arXiv:1303.6645v1 [astro-ph.EP] 26 Mar 2013

S-Type and P-Type Habitability in Stellar Binary Systems: A Comprehensive Approach
I. Method and Applications

M. Cuntz

Department of Physics

University of Texas at Arlington, Arlington, TX 76019-0059;

`cuntz@uta.edu`

ABSTRACT

A comprehensive approach is provided to the study of both S-type and P-type habitability in stellar binary systems, which in principle can also be expanded to systems of higher order. P-type orbits occur when the planet orbits both binary components, whereas in case of S-type orbits the planet orbits only one of the binary components with the second component considered a perturbator. The selected approach encapsulates a variety of different aspects, which include: (1) The consideration of a joint constraint including orbital stability and a habitable environment for a putative system planet through the stellar radiative energy fluxes ("radiative habitable zone"; RHZ) needs to be met. (2) The treatment of conservative, general and extended zones of habitability for the various systems as defined for the Solar System and beyond. (3) The providing of a combined formalism for the assessment of both S-type and P-type habitability; in particular, mathematical criteria are presented for which kind of system S-type and P-type habitability is realized. (4) Applications of the attained theoretical approach to standard (theoretical) main-sequence stars. In principle, five different cases of habitability are identified, which are: S-type and P-type habitability provided by the full extent of the RHZs; habitability, where the RHZs are truncated by the additional constraint of planetary orbital stability (referred to as ST and PT-type, respectively); and cases of no habitability at all. Regarding the treatment of planetary orbital stability, we utilize the formulae of Holman & Wiegert (1999) [AJ 117, 621], as also used by previous studies, although other methods for the assessment of stability / instability transitions in planetary systems as, e.g., the Lyapunov exponent criterion have also become available. In this work we focus on binary systems in circular orbits. Future applications will also consider binary systems in elliptical orbits and provide thorough comparisons to other methods and results given in the literature.

Subject headings: astrobiology — binaries: general — celestial mechanics — planetary systems

1. Introduction

Starting more than a decade ago considerable observational evidence has been obtained indicating that planets are able to exist in stellar binary (and higher order) systems; see results and discussions by, e.g., Patience et al. (2002), Eggenberger et al. (2004), and Eggenberger et al. (2007). These observations are in line with the empirical finding that binary (and higher order) systems occur in high frequency in the local Galactic neighborhood (Duquennoy & Mayor 1991; Lada 2006; Raghavan et al. 2006; Bonavita & Desidera 2007; Raghavan et al. 2010). For example, Raghavan et al. (2010) presented results of a detailed analysis of companions to solar-type stars, based on a sample size of 454, and concluded that the overall fractions of double and triple systems are about 33% and 8%, respectively, if all confirmed stellar and brown dwarf companions are accounted for.

The fairly frequent occurrence of planets in binary systems is furthermore consistent with the presence of debris disks in a considerably large number of main-sequence star binary systems (e.g., Trilling et al. 2007). In principle, as discussed by Perryman (2011), planets in binary systems can be identified through two different venues: First, binaries or multiple star systems can be surveyed for the presence of planets by utilizing the established detection methods. Second, stars with detected planets can be scrutinized afterward to check if they possess one or more widely separated stellar companion(s); in this case, the planet(s) will also be categorized as belonging to a binary (or higher order) system.

From the view point of orbital mechanics, there are two different kinds of possible orbits (notwithstanding positions near the Lagrangian points L_4 and L_5) for planets in binary systems: S-type and P-type orbits (Dvorak 1982). A P-type orbit is given when the planet orbits both binary components, whereas in case of an S-type orbit the planet orbits only one of the binary components with the second component behaving as a perturbator. Eggenberger et al. (2004) presented a list of 15 planet-bearing binary systems with all planets in S-type orbits. They constitute mostly wide binaries with separation distances of up to $\sim$6400 AU; however, smaller separation distances on the order of 20 AU or less have also been identified. In the meantime, systems with planets in P-type orbits have also been identified. Arguably, the most prominent case is Kepler-16, as reported by Doyle et al. (2011) and previously suggested by Slawson et al. (2011), containing a Saturnian mass circumbinary planet. Quarles et al. (2012) have subsequently studied this system regarding the possibility of habitable exoplanets and habitable exomoons. Recently, a transiting circumbinary multiplanet system, i.e., Kepler-47, has also been identified (Orosz et al. 2012).

There is a significant body of literature devoted to the study of habitability in binary systems as well as in multiplanetary systems, which often also encompass stellar evolutionary considerations. Examples include the work by Jones et al. (2001), Noble et al. (2002), Menou & Tabachnik (2003), Asghari et al. (2004), Sándor et al. (2007), Takeda et al. (2008), Dvorak et al. (2010), Jones & Sleep (2010), and Kopparapu & Barnes (2010). An important aspect that has received increased recognition in the literature is that in order for habitability to exist and to be maintained, a joint constraint that includes both orbital stability and a habitable environment for a system planet through the stellar radiative energy fluxes needs to be met. In the framework of this paper, the zone related to this latter requirement will subsequently be referred to as *radiative habitable zone* (RHZ), which constitutes a necessary, though often insufficient, condition for the existence of circumstellar habitability.

Previous work, mostly concentrated on the existence of habitability in single star multiplanetary systems, rendered the publication of detailed "stability catalogs" for the habitability zones of extrasolar planetary systems (e.g., Menou & Tabachnik 2003; Sándor et al. 2007; Takeda et al. 2008; Dvorak et al. 2010; Kopparapu & Barnes 2010). For example, Menou & Tabachnik (2003) quantified the dynamical habitability of 85 planetary systems by considering the perturbing influence of giant planets beyond the traditional Hill sphere for close encounters with the theoretical terrestrial planets. They concluded that a significant fraction of the identified extrasolar planetary systems are unable to harbor habitable terrestrial planets. A statistical study on the stability of Earth-mass planets orbiting solar-mass stars in presence of stellar companions focusing on both the statistical properties of ejection times and the general prospects of planetary habitability was given by Fatuzzo et al. (2006). Additional work providing stability assessments for various observed extrasolar planetary systems based on detailed stability maps was given by Sándor et al. (2007). Takeda et al. (2008) explored the orbital stability or planets in double-planet systems for binaries by supplying an analytic framework based on secular perturbation theory; they also provided dynamical classification categories. Additional stability analyses to assess the habitability of planetary systems based on detailed numerical simulations were given by Dvorak et al. (2010) and Kopparapu & Barnes (2010); note that the study of Dvorak et al. (2010) also dealt with a limited cases of planets in double star systems in orbit either around one stellar component (S-type) or around both components (S-type).

The study of planetary dynamics and habitable planet formation has meanwhile been described by, e.g., Quintana & Lissauer (2010) and Haghighipour et al. (2010). They show that Earth-mass planets are, in principle, able to form in stellar binary systems, also many of the physical, chemical, and astrobiological details are not fully understood, including relationships between the formed planets and the orbital and physical properties of their host stars. The overarching conclusion of those investigations is that habitable planets in stellar binary (and, as anticipated, in higher-order systems) are, in general, possible, which is a stark motivation for providing a comprehensive study of S-type and P-type habitability in binary systems. The approach adopted in this study will be entirely analytic. Specifically, it will consider both S-type and P-type habitable orbits in the view of the joint constraint including orbital stability and a habitable environment for a system planet through the appropriate amount of the stellar radiative energy fluxes. In an earlier study, Eggl et al. (2012) focused on S-type habitability in binary systems taking into account both circular and elliptical orbits for the stellar binary components; this latter aspect is however beyond the scope of the present work as we solely focus on systems in circular orbits. Numerical studies for P-type habitable environments with applications to Kepler-16, Kepler-34, Kepler-35 and Kepler-47 have given by Kane & Hinkel (2013).

Our paper is structured as follows: In Sect. 2, we comment on the adopted main-sequence star parameters and single star habitability. In Sect. 3, we introduce our theoretical approach suitable for stellar systems of the order of N, although our focus will be on binary systems. In this regard both S-type and P-type orbits will be examined, and detailed mathematical criteria for the existence of S-type and P-type RHZs will be derived. In Sect. 4, we consider the joint constraint of planetary orbital stability and the stellar radiative environment (i.e., RHZs) for the establishment of circumstellar habitability. Applications regarding S-type and P-type systems as well as the introduction of the habitability classifications S, P, ST, and PT are given in Sect. 5. In Sect. 6, we convey our summary and conclusions.

2. Stellar Parameters and Single Star Habitability

In this study, S-type and P-type habitability is investigated mostly pertaining to standard (i.e., theoretical) main-sequence stars. The adopted stellar parameters, which are the stellar effective temperatures $T_{\rm eff}$, the stellar radii R_* (which together allow to define the stellar luminosities L_*), and the stellar masses M_* are mainly based on the work by Gray (2005) (see his Table B.1) that assumes detailed photospheric spectral analyses. If a mass value in Gray's table appeared to be inaccurate due to numerical truncations, it was recalculated if more accurate values for the surface gravity log g_* and the stellar radii were given. For stellar spectral types with no data available, the missing data were computed by employing biparabolic interpolation.

The exception, however, are data for stars of spectral type K5 V and below. In this regard we relied on the results from the spectral models of R. L. Kurucz and collaborators.

They took into account hundreds of millions of spectral lines for a large set of atoms and molecules; see Castelli & Kurucz (2004) and Kurucz (2005) for details. The effective temperatures implied by these models are in close similarity to those given by Gray (2005) for most types of stars; however, Gray (2005) reports consistently higher effective temperatures for stars of spectral type late-K and M; for the latter, the difference amounts to nearly 300 K. Table 1 depicts the stellar parameters adopted for the present work.

An alternative approach expected to provide very similar results for either the stellar luminosity or the stellar mass (with the other parameter taken as fixed) is the employment of a mass–luminosity relationships applicable to main-sequence stars. The work by Reid (1987), as well as data from subsequent studies, yield

$$\frac{L_*}{L_\odot} \ = \ \eta \Bigl(\frac{M_*}{M_\odot} \Bigr)^\alpha \tag{1}$$

with $\eta = 0.23$ and $\alpha = 2.3$ for $M_* < 0.43 \ M_\odot$ and $\eta = 1$ and $\alpha = 4.0$ for $M_* \geq 0.43 \ M_\odot$. At the high-mass end, this relationship holds until about $M_* = 2 \ M_\odot$. It also becomes increasingly inaccurate for low-mass M dwarfs. Fortunately, the domains of applicability for Eq. (1) is consistent with most studies of binary habitability; see, e.g., Eggl et al. (2012) as example.

Next we focus on single star habitability, i.e., the evaluation of various limits of habitable zones (HZs), which in the solar case shall be referred to as s_ℓ. Previous work by, e.g., Kasting et al. (1993) distinguished between conservative (CHZ) and the generalized habitable zone (GHZ), which can also be evaluated for general main-sequence stars, and other types of stars as well. For the Sun, the limits of the CHZ are given as 0.95 and 1.37 AU ($\ell = 2$ and 4, respectively), whereas for the GHZ, they are given as 0.84 and 1.67 AU ($\ell = 1$ and 5, respectively); see Table 2.

The physical significance of the various kinds of HZs obtained by Kasting et al. (1993) can be summarized as follows: The GHZ is defined as bordered by the runaway greenhouse effect (inner limit) and the maximum greenhouse effect (outer limit). Concerning the latter it is assumed that a cloud-free CO_2 atmosphere is still able to provide a surface temperature of 273 K. The inner limit of the CHZ is defined by the onset of water loss. In this case, a wet stratosphere is assumed to exist where water is lost by photodissociation and subsequent hydrogen escape to space. Furthermore, the outer limit of the CHZ is defined by the first CO_2 condensation attained by the onset of formation of CO_2 clouds at a temperature of 273 K; see, e.g., Underwood et al. (2003) and Selsis et al. (2007) for further details. Table 3 conveys the results for the HZs of the different type of main-sequence stars considered in the present study.

For the outer edge of circumstellar habitability, even less stringent limits have been introduced in the meantime (e.g., Forget & Pierrehumbert 1997; Mischna et al. 2000). They are based on the assumption of relatively thick planetary CO_2 atmospheres as well as strong backwarming that may further be enhanced by CO_2 crystals and clouds. These limits, which in case of the Sun correspond to 2.4 AU ($s_\ell = s_6$), conform to the extended habitable zone (EHZ), have also been taken into account in our study, although the significance of the EHZ has meanwhile been criticized as a result of detailed planetary radiative transfer models (Halevy et al. 2009). Moreover, in the framework of the present study, we also consider planetary Earth-equivalent positions defined as $R_{\oplus,\mathrm{eqv}} \simeq \sqrt{L_*/L_\odot}$ and labelled as $s_\ell = s_3$; see Table 3. It is meant as an intriguing reference distance of habitability both regarding single stars and stellar binary systems.

3. Theoretical Approach

3.1. Basic Equations

Next we introduce the governing equations for investigating the RHZs of binary systems pertaining to both S-type and P-type orbits. This approach targets the requirement of providing a habitable environment for a system planet based on the radiative energy fluxes of the stellar components. The requirement of planetary orbital stability will be disregarded for now; it will be revisited in Sect. 4. The importance of orbital stability for allowing circumstellar habitability in stellar binaries will, however, be considered in an appropriate and consistent manner in the main body of the study.

For a star of luminosity L_i, given in units of solar luminosity $L_\odot$, the distance d_i of the habitability limit s_ℓ as identified for the Sun, which may constitute either an inner or outer limit of habitability (except $\ell = 3$), is given as

$$d_i \ = \ s_\ell \sqrt{\frac{L_i}{S_{\mathrm{rel},i} L_\odot}} \tag{2}$$

In case of a multiple star system of order N with distances d_i, the limit of habitability related to s_ℓ is given as[1]

$$\sum_{i=1}^{N} \frac{L_i}{S_{\mathrm{rel},i} d_i^2} \ = \ \frac{L_\odot}{s_\ell^2} \ . \tag{3}$$

---

[1] This equation is analogous to an equivalent equation of electrostatics relating a general distribution of charges to the resulting electrostatic potential in free space (Jackson 1999, see p. 40, Eq. (1.48)); a modified version of Eq. (3) has previously been considered by, e.g., Eggl et al. (2012).

In Eq. (2) and (3), $S_{\mathrm{rel},i} = S_{\mathrm{rel},i}(T_{\mathrm{eff}})$ describes the stellar flux in units of the solar constant that is a function of the stellar effective temperature T_{eff} (e.g., Kasting et al. 1993; Underwood et al. 2003). Specifically, using the formalism[2] by Selsis et al. (2007), we find that

$$S_{\mathrm{rel},i} \;=\; \Big(\frac{s_\ell}{s_\ell - a_z T_* - b_z T_*^2}\Big)^2 \tag{4}$$

with s_ℓ, a_z, and b_z in AU, and $T_* = T_{\mathrm{eff}} - 5700$ in K. Selsis et al. (2007) also found that for $s_\ell < 1$, corresponding to inner limits of habitability, the fitting parameters are given as $a_z = 2.7619 \times 10^{-5}$ and $b_z = 3.8095 \times 10^{-9}$, whereas for $s_\ell > 1$, corresponding to outer limits of habitability, they are given as $a_z = 1.3786 \times 10^{-4}$ and $b_z = 1.4286 \times 10^{-9}$; note that $s_\ell \equiv 1$ corresponds to the customary notion of Earth-equivalent positions. Appropriate values for s_ℓ are given in Table 2.

In the following, we will focus on the case of binary systems, i.e., $N = 2$. In this case Eq. (3) reads

$$\frac{L_1}{S_{\mathrm{rel},1} d_1^2} + \frac{L_2}{S_{\mathrm{rel},2} d_2^2} \;=\; \frac{L_\odot}{s_\ell^2} \tag{5}$$

with

$$d_1^2 \;=\; a^2 + z^2 + 2az\cos\varphi \tag{6a}$$

$$d_2^2 \;=\; a^2 + z^2 - 2az\cos\varphi\ . \tag{6b}$$

Here a denotes the semidistance of binary separation, z the distance of a position at the habitability limit contour (which later on will be referred to as "radiative habitable limit", see below), and φ the associated angle; see Fig. 1 for information on the coordinate set-up for both S-type and P-type orbits. We will also assume $L_1 \geq L_2$ without loss of generality.

With L_i' defined as

$$L_i' = \frac{L_i}{L_\odot S_{\mathrm{rel},i}}, \tag{7}$$

---

[2] Here $S_{\mathrm{rel},i}$ represents the normalized stellar flux in units of the solar constant, 1368 W m^{-2}, given by the stellar spectral energy distribution. Therefore, ordinarily, no s_ℓ dependence for $S_{\mathrm{rel},i}$ should exist. However, the formulae by Selsis et al. (2007) utilize previous results by Kasting et al. (1993) who provided numerical values for limits of habitability for different types of stars considering various limit definitions (i.e., s_ℓ values identified for the Sun). But Kasting et al. (1993) used for the solar effective temperature an unusually low value of 5700 K instead of 5777 K as currently accepted (e.g., Stix 2004). Hence transforming the polynomial fit based on the work by Kasting et al. aimed at considering the correct solar effective temperature renders a weak dependence on s_ℓ for the $S_{\mathrm{rel},i}$ values. In contrast the method by Underwood et al. (2003) provides a polynomial fit for $S_{\mathrm{rel},i}$ without considering the solar temperature revision. An alternative method has been used by Cuntz & Yeager (2009) and subsequent work. In this approach the polynomial fit by Underwood et al. (2003) is corrected via a triangular function based on data for stars of spectral type F0 V, G0 V, and K0 V. As a result the corresponding $S_{\mathrm{rel},i}$ values do also not show any s_ℓ dependence.

henceforth referred to as *recast stellar luminosity*, $z(\varphi)$ is given as

$$z^4 + A_2 z^2 + A_1 z + A_0 \ = \ 0 \tag{8}$$

with

$$\begin{aligned} A_2 &= 2a^2(1-2\cos^2\varphi) - s_\ell^2(L_1' + L_2') && \text{(9a)} \\ A_1 &= 2as_\ell^2 \cos\varphi(L_1' - L_2') && \text{(9b)} \\ A_0 &= a^4 - a^2 s_\ell^2(L_1' + L_2') \ . && \text{(9c)} \end{aligned}$$

Equation (8) constitutes a fourth-order algebraic equation that is known to possess four possible solutions (Bronshtein & Semendyayev 1997), although some (or all) of them may constitute unphysical solutions, i.e., $z(\varphi)$ having a complex or imaginary value. The adopted coordinate system constitutes, in essence, a polar coordinate system except that negative values for z are permitted; in this case the position of z is found on the opposite side of angle φ.

In principle, it is possible to consider for Eq. (8) to only have solutions for $z(\varphi)$ given as $z \geq 0$; in this case the entire interval for $z(\varphi)$, which is $0 \leq \varphi < 2\pi$, needs to be examined. The following types of solutions are identified: For S-type orbits, two solutions exist in the intervals centered at $\varphi = 0$ and at $\varphi = \pi$ (or one coinciding solution at each tangential point); see Fig. 1. However, there will be no solution in the typically relatively large intervals containing $\varphi = \pi/2$ and $\varphi = 3\pi/2$. Clearly, the size of any of those intervals critically depends on the system parameters a, L_1', L_2', and s_ℓ, as expected. For P-type orbits, on the other hand, there will be one solution for each value of φ in the range of $0 \leq \varphi < 2\pi$.

However, in general, negative values for the solutions of $z(\varphi)$ also exist. If taken into account, it will be sufficient to restrict the evaluation of Eq. (8) to the range $0 \leq \varphi \leq \pi$. In this case, for S-type orbits, there will be four solutions in the intervals with endpoints $\varphi = 0$ and $\varphi = \pi$, as well as two solutions (if $L_1' \neq L_2'$) in a more extended interval containing these points. Also, a pair of solutions will become one coinciding solution at each tangential point. However, again, there will be no solution in the interval containing $\varphi = \pi/2$. In case of P-type orbits, there will be two solutions for any value of φ in the range of $0 \leq \varphi \leq \pi$. We will revisit this assessment in conjunction with the algebraic method for attaining the solution z_i; additionally, detailed mathematical criteria will be given for the existence of RHZ for S-type and P-type orbits.

Next we will focus on equal-star binary systems. Detailed solutions for general binary systems (i.e., systems of stellar components with by default unequal masses, luminosities, and effective temperatures) pertaining to both S-type and P-type orbits will be given in Sect. 3.3. Both subsections will be aimed at deriving RHZs; see, e.g., Williams & Pollard (2002) for general discussions on the role of RHZs for the attainment of habitability in star–planet systems. However, strictly speaking, they will deal with identifying *radiative habitable limits* (RHLs) connected to a distinct value of s_ℓ noting that manifesting a RHZ requires that the RHL for $s_{\ell,\mathrm{out}}$ to be located *completely outside* of the RHL for $s_{\ell,\mathrm{in}}$ with $s_{\ell,\mathrm{in}}$ and $s_{\ell,\mathrm{out}}$ appropriately paired. A summary about the existence and structure of the RHZs, encompassing the radiative CHZs, GHZs, and EHZs, will be given in Sect. 3.4; this subsection will also convey cases where no RHZs exist due to the behavior of the RHLs regarding $s_{\ell,\mathrm{in}}$ and $s_{\ell,\mathrm{out}}$.

However, in none of these subsections the requirement of orbital stability for possible system planets is considered, which in general entails further restrictions on the manifestation of binary system habitability. This latter aspect will be discussed in Sect. 4 and considered in detail in the subsequent set of applications (see Sect. 5).

3.2. Equal-Star Binary Systems

Now we focus on the special case of equal-star binary systems, i.e., stars of identical recast luminosities, i.e., $L'_1 = L'_2 = L'$. For theoretical main-sequence stars this assumption also implies $S_{\mathrm{rel},1} = S_{\mathrm{rel},2}$ and $M_1 = M_2$; this latter assumption about the stellar masses is relevant for the orbital stability constraint of system planets. With $A_1 = 0$, Eq. (8) now constitutes a biquadratic equation that can be solved in a straightforward manner. The other coefficients are given as

$$\begin{aligned} A_2 &= 2a^2(1-2\cos^2\varphi) - 2s_\ell^2 L' && \text{(10a)} \\ A_0 &= a^4 - 2a^2 s_\ell^2 L' \ . && \text{(10b)} \end{aligned}$$

Thus, the solution of Eq. (8) is given as

$$z \ = \ \pm\sqrt{-a^2(1-2\cos^2\varphi) + s_\ell^2 L' \pm \sqrt{D_2}} \qquad (11)$$

with

$$D_2 \ = \ s_\ell^4 L'^2 + 4a^4\cos^4\varphi - 4a^2\cos^2\varphi(a^2 - s_\ell^2 L'). \qquad (12)$$

With known systems parameters, which are a, s_ℓ, and L', the function $z(\varphi)$, describing the habitability limits for the binary system associated with inner limit and outer limit values s_ℓ derived for the Sun (see Sect. 2 for details) can be obtained in a straightforward manner.

Owing to the system symmetry, the existence of S-type and P-type RHLs can be found by attaining the solutions of Eq. (11) for $\varphi = 0$ and $\varphi = \pi/2$. First we examine the solutions of Eq. (11) for $\varphi = 0$, i.e., $\cos\varphi = 1$, which are given as

$$z \; = \; \pm\sqrt{a^2 + s_\ell^2 L' \pm s_\ell \sqrt{s_\ell^2 {L'}^2 + 4a^2 L'}} \; . \tag{13}$$

The total number of solutions for z_i is four as expected; it collapses to two solutions if the value of the interior square root approaches zero, which obviously reflects the mathematical limit for the existence of S-type RHLs. Thus, S-type RHLs exist if

$$a \; \geq \; s_\ell \sqrt{2L'} \; . \tag{14}$$

Next we examine the solutions of Eq. (11) for $\varphi = \pi/2$, i.e., $\cos\varphi = 0$. This allows us to gauge the existence of P-type RHLs; the latter implies two solutions of Eq. (11) regardless of the value for φ. Here the solution is given as

$$z \; = \; \pm\sqrt{-a^2 + 2s_\ell^2 L'} \; . \tag{15}$$

Thus, the condition for P-type RHLs is given as

$$a \; \leq \; s_\ell \sqrt{2L'} \; . \tag{16}$$

Therefore, Eqs. (14) and (16) allow to identify conditions for S-type and P-type RHLs, respectively, for equal-star binary systems, which depend on the systems parameters a, s_ℓ, and L'. Comparing Eqs. (14) and (16) also implies that the joint existence S-type and P-type habitability in equal-star binary systems is impossible, irrespectively of the planetary orbital stability requirement (see Sect. 4), noting that the latter imposes an additional constraint on habitability even when the RHZ-related conditions are met. Figure 2 depicts the borders of the S-type and P-type radiative habitable limits, i.e., RHLs, for different values of s_ℓ in regard to a and L'.

3.3. General Binary Systems

3.3.1. Method of Solution

We now focus on obtaining solutions for our key equation, Eq. (8), pertaining to S-type and P-type RHLs for general binary systems, i.e., $L_1' \neq L_2'$. Following, e.g., Abramowitz & Stegun (1972) and Beyer (1987), the set of solutions for a fourth-order polynomial reads

$$
\begin{aligned}
z_1 &= -\frac{1}{2}\mathcal{C} - \frac{1}{2}\mathcal{D} && \text{(17a)}\\
z_2 &= -\frac{1}{2}\mathcal{C} + \frac{1}{2}\mathcal{D} && \text{(17b)}\\
z_3 &= +\frac{1}{2}\mathcal{C} - \frac{1}{2}\mathcal{E} && \text{(17c)}\\
z_4 &= +\frac{1}{2}\mathcal{C} + \frac{1}{2}\mathcal{E} && \text{(17d)}
\end{aligned}
$$

with[3]

$$
\begin{aligned}
\mathcal{C} &= \sqrt{-2a^2(1-2\cos^2\varphi) + s_\ell^2(L_1'+L_2') + y_1} && \text{(18a)}\\
\mathcal{D} &= \sqrt{s_\ell^2(L_1'+L_2') + 4as_\ell^2(L_1'-L_2')\mathcal{C}^{-1}\cos\varphi - 4a^2(1-2\cos^2\varphi) - y_1} && \text{(18b)}\\
\mathcal{E} &= \sqrt{s_\ell^2(L_1'+L_2') - 4as_\ell^2(L_1'-L_2')\mathcal{C}^{-1}\cos\varphi - 4a^2(1-2\cos^2\varphi) - y_1} && \text{(18c)}
\end{aligned}
$$

with y_1 as a solution of the resolvent cubic equation

$$
y^3 - A_2y^2 - A_0y + (4A_2A_0 - A_1^2) = 0 \tag{19}
$$

with A_0, A_1, and A_2 given by Eqs. (9a) to (9c). The solutions z_i given through Eqs. (17a) to (17d), if existing, are ordered as $z_1 < z_2 < z_3 < z_4$ for $\varphi = 0$, as well as for any other value of φ as subsequently identified. Although Eq. (19) has three possible solutions, there is only one appropriate choice for y_1 because it is necessary to avoid that all z_i obtained through Eq. (8) are of imaginary or conjugate complex value in cases where S-type or P-type RHLs exist.

The adopted method also requires a solution for y; see Eq. (19). The only acceptable solution for y, customarily denoted as y_1, is given as

$$
y_1 = -\frac{1}{3}\hat{a}_2 + (S+T) \tag{20}
$$

---

[3] In Eqs. (18b) and (18c) we also correct a typo in the sign preceeding the term of $\mathcal{C}^{-1}$.

with the substitutions

$$\begin{aligned} S &= \sqrt[3]{R+\sqrt{D_3}} && \text{(21a)} \\ T &= \sqrt[3]{R-\sqrt{D_3}} && \text{(21b)} \\ D_3 &= Q^3+R^2 && \text{(21c)} \end{aligned}$$

and with Q and R given as

$$\begin{aligned} Q &= \frac{1}{3}\hat{a}_1 - \frac{1}{9}\hat{a}_2^2 && \text{(22a)} \\ R &= -\frac{1}{2}\hat{a}_0 + \frac{1}{6}\hat{a}_1\hat{a}_2 - \frac{1}{27}\hat{a}_2^3 && \text{(22b)} \end{aligned}$$

while noting that

$$\begin{aligned} \hat{a}_0 &= 4A_0A_2 - A_1^2 && \text{(23a)} \\ \hat{a}_1 &= -4A_0 && \text{(23b)} \\ \hat{a}_2 &= -A_2\ . && \text{(23c)} \end{aligned}$$

These sets of equation can be solved and appropriate values for $z(\varphi)$ can be obtained. The results will depend on the system parameters a, L_1', L_2', and s_ℓ, as expected.

Next we describe the solutions for S-type and P-type RHLs in more detail. It is important to recognize that apriori choices about the existence of S-type and P-type RHLs are neither necessary nor possible as the existence of any of those RHLs is determined by the fulfillment of well-defined mathematical conditions; they will also be given in the following.

3.3.2. S-type Orbits

An analysis of the possible solutions for Eqs. (17a) to (17d) shows that for S-type RHLs valid solutions are obtained based on

$$\begin{aligned} S &= \sqrt[6]{R^2+K^2}\ (\cos\xi + i\sin\xi) && \text{(24a)} \\ T &= \sqrt[6]{R^2+K^2}\ (\cos\xi - i\sin\xi) && \text{(24b)} \end{aligned}$$

with

$$\begin{aligned} K &= \sqrt{|D_3|} && \text{(25a)} \\ \xi &= \frac{1}{3}\arctan\left(\frac{K}{R}\right) && \text{(25b)} \end{aligned}$$

with R given by Eq. (22b). Therefore, the solution of the resolvent cubic equation, Eq. (19), is given as

$$y_1 \;=\; -\frac{1}{3}\hat{a}_2 + 2\sqrt[6]{R^2+K^2}\cos\xi \tag{26}$$

For the values of z_i it is found that for the interval centered at $\varphi = 0$, z_1 and z_2 exhibit negative values, whereas z_3 and z_4 exhibit positive values with $z_1 < z_2 < z_3 < z_4$. Thus, z_1 and z_2 describe the RHL regarding star S1, whereas z_3 and z_4 describe the RHL regarding star S2 (see Fig. 1). Conversely, for the interval centered at $\varphi = \pi$, z_1 and z_2 again exhibit negative values and z_3 and z_4 exhibit positive values. In this case, z_1 and z_2 describe the RHL for star S2, whereas z_3 and z_4 describe the RHL for star S1. Surely, no solutions are obtained in the vicinities of $\varphi = \pi/2$ and $\varphi = 3\pi/2$, as expected. Thus, for each angle φ in the range of $0 \le \varphi < 2\pi$ the appropriate number of solutions is attained to describe S-type RHLs. However, due to symmetry, solutions are only needed for $0 \le \varphi \le \pi$.

The existence of S-type RHLs requires that $z_2 \le z_3$ because otherwise the two distinct S-type RHL contours about the two binary components would not be separated, which corresponds to the condition

$$\mathcal{C} \;\ge\; \frac{1}{2}(\mathcal{D}+\mathcal{E}) \ . \tag{27}$$

This equation can most likely be appropriately rewritten to provide an expression only containing the system parameters A_0, A_1, and A_2 defined through Eqs. (9a) to (9c). However, the resulting expression is expected to be highly intricate.

3.3.3. P-type Orbits

An analysis of the possible solutions for Eqs. (17a) to (17d) also shows that for P-type RHLs valid solutions require

$$D_3 \;\ge\; 0 \ ; \tag{28}$$

see Eq. (21c). The detailed evaluation of this condition requires the evaluation of various sets of equations denoted as Eqs. (9), (21), and (22); see Sect. 3.1 and 3.3.1.

In terms of the solutions for P-type RHLs it is found that for $0 \le \varphi < \pi/2$ and $3\pi/2 < \varphi \le 2\pi$, z_1 exhibit negative values and z_2 exhibit positive values, whereas z_3 and z_4 are undefined; they are also not needed for outlining P-type RHLs. Moreover, for the range of $\pi/2 < \varphi < 3\pi/2$, z_3 exhibit negative values and z_4 exhibit positive values, noting that z_1 and z_2 remain undefined. For $\pi/2$ and $3\pi/2$, removable singularities are identified, which can easily be fixed through interpolation taking values of z for neighboring angles of φ. In summary, for each angle φ in the range of $0 \le \varphi < 2\pi$ two values of z_i (i.e., one positive and one negative value) are identified allowing to determine P-type RHLs. However, due to symmetry, solutions are only needed for $0 \leq \varphi \leq \pi$.

Furthermore, through analytical transformations it can be shown that the condition depicted as Eq. (29) can be rewritten as

$$16 A_0 A_2^4 + 144 A_0 A_1^2 A_2 - 128 A_0^2 A_2^2 + 256 A_0^3 - 4 A_1^2 A_2^3 - 27 A_1^4 \ \leq \ 0 \ . \tag{29}$$

with A_0, A_1, and A_2 defined through Eqs. (9a) to (9c). A key aspect concerns the following: For P-type RHLs to exist for the system parameters a, L_1', L_2', and s_ℓ, it is necessary that the relations (28) and (29), which are equivalent, must be fulfilled for any angle of φ.

A subsequent evaluation of Eq. (29) shows that in case of equal-star binary systems, corresponding to $A_1 = 0$, Eq. (29) can be simplified as

$$(A_2^2 - 4A_0)^2 \geq 0 \ ; \tag{30}$$

this relationship is fulfilled in a trivial manner. However, if the positive root of Eq. (30) is taken, the then recovered relation between A_0 and A_2 leads to the relationship of Eq. (16) valid for the special case of equal-star binary systems previously discussed (see Sect. 3.2).

3.4. Calculation of RHZs for Binary Systems

The identification of the RHZs in binary systems requires the calculation of limits of habitability, i.e., RHLs, as discussed in Sect. 3.1. They need to be established for values of s_ℓ with $\ell = 1$, 2, 4, 5, and 6 (see Sect. 2 and Table 2), which are informed by model-dependent physical limits of habitability for the solar environment (e.g., Kasting et al. 1993). As part of the process, the parameters of s_ℓ need to be appropriately paired in terms of the inner and outer limits of habitability. For the CHZ the parameters $(s_{\ell,\mathrm{in}}, s_{\ell,\mathrm{out}})$ need to be paired as (s_2, s_4), whereas for the GHZ, they need to paired as (s_1, s_5). For the EHZ, the parameters of s_ℓ need to paired as (s_1, s_6), considering that both the CHZ and the GHZ shall be viewed as subdomains of the EHZ.

For S-type and P-type orbits, the radiative zones of habitability RHZ(z), which constitutes a circular region (annulus) around each star S1 and S2 (S-type) or both stars (P-type) can be determined as

$$\mathrm{RHZ}(z) \ = \ \mathrm{Min}\Big(\mathcal{R}(z,\alpha)\Big)\Big|_{s_{\ell,\mathrm{out}}} - \mathrm{Max}\Big(\mathcal{R}(z,\alpha)\Big)\Big|_{s_{\ell,\mathrm{in}}} \tag{31}$$

and

$$\mathrm{RHZ}(z) \ = \ \mathrm{Min}\Big(\mathcal{R}(z,\varphi)\Big)\Big|_{s_{\ell,\mathrm{out}}} - \mathrm{Max}\Big(\mathcal{R}(z,\varphi)\Big)\Big|_{s_{\ell,\mathrm{in}}} , \tag{32}$$

respectively; see Fig. 1 for coordinate information. Here $\mathcal{R}(z,\alpha)$ and $\mathcal{R}(z,\varphi)$ describe the areas bordered by the RHLs defined by $s_{\ell,\mathrm{in}}$ and $s_{\ell,\mathrm{out}}$. The calculation of the extrema is applied to the angles α and φ for the intervals $0 \le \alpha \le \pi$ and $0 \le \varphi \le \pi/2$, respectively; note that we assumed $L_1 \ge L_2$ without loss of generality. In the S-type case the calculation of the extrema pertaining to the RHZ values is based on the angular coordinate α instead of φ; however, the angular coordinate φ is still needed for the calculation of z_i as part of the overall approach toward identifying S-type and P-type habitability.

Figure 3 depicts examples of RHLs and RHZs for different types of systems. In the S-type case the RHLs are bended toward the center of the system, whereas in the P-type case they are of notable elliptical shape. The RHZs always constitute circular annuli obtained through inspecting the appropriate minima and maxima of the RHLs. The examples as depicted include S-type and P-type systems with separation distances ($2a$) of 0.5 AU and 5.0 AU, respectively. Cases of both equal-star and non-equal star binaries are selected. The focus of this figure is the identification of the appropriate circular region (annulus) for each case. The figure also indicates the portions within the $\mathcal{R}(z,\alpha)$ and $\mathcal{R}(z,\varphi)$ domains that are not part of the RHZ(z) annuli.

Next we determine the values for the extrema pertaining to RHZ(z) following Eqs. (31) and (32) based on the solutions of Eq. (8) given as Eqs. (17a) to (17d). Noting that for existing values of z_i they are ordered as $z_1 < z_2 < z_3 < z_4$, and negative values for z_i are permitted it is sufficient for S-type orbits, both for the stars S1 and S2, to only consider solutions for $\varphi = 0$. For P-type orbits, a more detailed assessment is required (see below). For S-type orbits, regarding star S1, the extrema are obtained as

$$\mathrm{RHZ_{in}} = \mathrm{Max}(\mathcal{R}(z,\alpha))\Big|_{s_{\ell,\mathrm{in}}} = |a + z_2(0)|\Big|_{s_{\ell,\mathrm{in}}} \tag{33a}$$

$$\mathrm{RHZ_{out}} = \mathrm{Min}(\mathcal{R}(z,\alpha))\Big|_{s_{\ell,\mathrm{out}}} = |a + z_1(0)|\Big|_{s_{\ell,\mathrm{out}}} , \tag{33b}$$

and for star S2, they are obtained as

$$\mathrm{RHZ_{in}} = \mathrm{Max}(\mathcal{R}(z,\alpha))\Big|_{s_{\ell,\mathrm{in}}} = |a - z_3(0)|\Big|_{s_{\ell,\mathrm{in}}} \tag{34a}$$

$$\mathrm{RHZ_{out}} = \mathrm{Min}(\mathcal{R}(z,\alpha))\Big|_{s_{\ell,\mathrm{out}}} = |a - z_4(0)|\Big|_{s_{\ell,\mathrm{out}}} . \tag{34b}$$

The size of each annulus Δ_{RHZ} for pairs $(s_{\ell,\mathrm{in}}, s_{\ell,\mathrm{out}})$ is given as $\Delta_{\mathrm{RHZ}} = \mathrm{RHZ_{out}} - \mathrm{RHZ_{in}}$.

For P-type orbits, the extrema are given as follows

$$\mathrm{RHZ_{in}} = \mathrm{Max}(\mathcal{R}(z,\varphi))\Big|_{s_{\ell,\mathrm{in}}} = |z_1(0)|\Big|_{s_{\ell,\mathrm{in}}} \tag{35a}$$

$$\mathrm{RHZ_{out}} = \mathrm{Min}(\mathcal{R}(z,\varphi))\Big|_{s_{\ell,\mathrm{out}}} = |z_1(\varphi_{\mathrm{out}})|\Big|_{s_{\ell,\mathrm{out}}} , \tag{35b}$$

We note that for the angle $\varphi_{\rm out}$ for RHZ$_{\rm out}$ no straightforward expression[4] exists; it is located in the interval $0 < \varphi_{\rm out} \le \pi/2$. But it can be found numerically as it is given by the angle where the minimum of $|z(\varphi)|$ occurs.

There is also another complication in the identification of RHZ P-type orbits. Generally, it is required for the RHL of $s_{\ell,{\rm out}}$ to be located completely outside of the RHL of $s_{\ell,{\rm in}}$, i.e.,

$$\mathrm{Min}\Big(\mathcal{R}(z,\varphi)\Big)\Big|_{s_{\ell,{\rm out}}} \ \ge \ \mathrm{Max}\Big(\mathcal{R}(z,\varphi)\Big)\Big|_{s_{\ell,{\rm in}}} \ . \tag{36}$$

However, this condition is violated in some models, especially for relatively large values of a as well as relatively small ratios of L'_2/L'_1. In this case the RHZ for $(s_{\ell,{\rm in}}, s_{\ell,{\rm out}})$ is nullified, a behavior that may occur for the pairings (s_2, s_4), (s_1, s_5), and (s_1, s_6), corresponding to the CHZ, GHZ, and EHZ, respectively. In this regard the existence of the CHZ is in most jeopardy as $(s_4 - s_2)$ constitutes the smallest bracket among the various kinds of HZs (see Table 2). Also note that if the GHZ is nullified, the CHZ will be nullified as well considering that the CHZ (if existing) is entirely located within the GHZ. Likewise, if the EHZ is nullified the existence of both the CHZ and the GHZ will be nullified. Detailed examples will be given in the application segment of this paper; see Sect. 5.2 and 5.3 for details. However, this phenomenon does not exist for RHZ S-type orbits. The reason is that in comparable settings the RHL for $s_{\ell,{\rm out}}$ is mathematically undefined, which also amounts to a limitation for the extent of the associated RHZ.

For equal-star binary systems, with the property of $L'_1 = L'_2$, expressions for RHZ$_{\rm in}$ and RHZ$_{\rm out}$ for S-type and P-type orbits can be obtained based on Eqs. (13) and (15). For S-type orbits we find

$$\mathrm{RHZ}_{\rm in} \ = \ \left| a - \sqrt{a^2 + s_{\ell,{\rm in}}^2 L' + s_{\ell,{\rm in}} \sqrt{s_{\ell,{\rm in}}^2 {L'}^2 + 4a^2 L'}} \right| \tag{37a}$$

$$\mathrm{RHZ}_{\rm out} \ = \ \left| a - \sqrt{a^2 + s_{\ell,{\rm out}}^2 L' - s_{\ell,{\rm out}} \sqrt{s_{\ell,{\rm out}}^2 {L'}^2 + 4a^2 L'}} \right| \ . \tag{37b}$$

It is furthermore intriguing to explore the limit $a \gg s_{\ell,{\rm out}}\sqrt{L'}$, noting that $s_{\ell,{\rm out}}\sqrt{L'} > s_{\ell,{\rm in}}\sqrt{L'}$. In this case it is found that

$$\mathrm{RHZ}_{\rm in} \ = \ \left| \frac{1}{2a} \Big(s_{\ell,{\rm in}}^2 L' + s_{\ell,{\rm in}} \sqrt{s_{\ell,{\rm in}}^2 {L'}^2 + 4a^2 L'} \Big) \right| \tag{38a}$$

$$\mathrm{RHZ}_{\rm out} \ = \ \left| \frac{1}{2a} \Big(s_{\ell,{\rm out}}^2 L' - s_{\ell,{\rm out}} \sqrt{s_{\ell,{\rm out}}^2 {L'}^2 + 4a^2 L'} \Big) \right| \ . \tag{38b}$$

---

[4] Due to the nature of the underlying equations for z_i, an analytic expression for φ is deemed possible. However, it is expected to be highly complicated and thus left as a student exercise.

Moreover, in the limit of $a \to \infty$, the expressions for single star HZs regarding $\mathrm{RHZ_{in}}$ and $\mathrm{RHZ_{out}}$ are recovered, as expected, which are given as $s_{\ell,\mathrm{in}}\sqrt{L'}$ and $s_{\ell,\mathrm{out}}\sqrt{L'}$, respectively. They are in agreement with the expressions previously obtained by Kasting et al. (1993), Underwood et al. (2003), Selsis et al. (2007), and others.

Results for P-type orbits can be obtained considering

$$\begin{aligned}
\mathrm{RHZ_{in}} &= \mathrm{Max}\Big(\mathcal{R}(z,0), \mathcal{R}(z,\frac{\pi}{2})\Big)\Big|_{s_{\ell,\mathrm{in}}} = \mathcal{R}(z,0)\Big|_{s_{\ell,\mathrm{in}}} && (39a)\\
\mathrm{RHZ_{out}} &= \mathrm{Min}\Big(\mathcal{R}(z,0), \mathcal{R}(z,\frac{\pi}{2})\Big)\Big|_{s_{\ell,\mathrm{out}}} = \mathcal{R}(z,\frac{\pi}{2})\Big|_{s_{\ell,\mathrm{out}}} \ . && (39b)
\end{aligned}$$

In this case we find

$$\begin{aligned}
\mathrm{RHZ_{in}} &= \sqrt{a^2 + s_{\ell,\mathrm{in}}^2 L' + s_{\ell,\mathrm{in}}\sqrt{s_{\ell,\mathrm{in}}^2 L'^2 + 4a^2 L'}} && (40a)\\
\mathrm{RHZ_{out}} &= \sqrt{-a^2 + 2 s_{\ell,\mathrm{out}}^2 L'} && (40b)
\end{aligned}$$

based on the system parameters a, L', $s_{\ell,\mathrm{in}}$, and $s_{\ell,\mathrm{out}}$. Additionally, the requirement to avoid that the RHL for $s_{\ell,\mathrm{out}}$ to be partially or completely located inside of the RHL for $s_{\ell,\mathrm{in}}$, see Eq. (36), entails

$$a^2 + s_{\ell,\mathrm{in}}\sqrt{a^2 L' + \frac{1}{4} s_{\ell,\mathrm{in}}^2 L'^2} \ \leq \ -\frac{1}{2} s_{\ell,\mathrm{in}}^2 L' + s_{\ell,\mathrm{out}}^2 L' \ , \qquad (41)$$

which allows to set constraints on the separation distance $2a$ of the binary system noting that the values of L', $s_{\ell,\mathrm{in}}$, and $s_{\ell,\mathrm{out}}$ are all subject to distinct restrictions, particularly in case of main-sequence stars (see Tables 1 and 2). Depictions of the condition (41) for the pairings (s_2, s_4), (s_1, s_5), and (s_1, s_6), corresponding to the CHZ, GHZ, and EHZ, respectively, are given in Fig. 4. A similar expression is expected to hold for nonequal-star binaries, though it will be highly complicated.

4. Constraints on Habitability due to Planetary Orbital Stability

A primary constraint on planetary habitability is that planets are required to exist in the HZ for a sufficient amount of time allowing basic forms of life to emerge and develop. In order to adhere to this criterion, planetary orbital stability is required. There is a significant body of literature devoted to this topic, including studies of binary and multi-planetary systems, which often also consider aspects of stellar evolution (e.g., Jones et al. 2001; Noble et al. 2002; Menou & Tabachnik 2003; Sándor et al. 2007; Takeda et al. 2008; Dvorak et al. 2010; Haghighipour et al. 2010; Kopparapu & Barnes 2010).

Early studies of planetary orbital stability pertaining to planets in both S-type and P-type orbits demonstrated that planets can exist in systems of binary stars for 3000 binary periods Dvorak (1984, 1986). Although these investigations considered relatively short integration times, Dvorak determined upper and lower bounds of planetary orbital stability considering the orbital elements, semimajor axis and eccentricity, of the proposed binary stars. Since this pioneering work, many additional studies have been performed. The foremost investigation extended the original study by a factor of 10 in integration times and an extended range of orbital elements (Holman & Wiegert 1999). In addition, the nature of the bounding formula was derived and discussed using a more statistical framework. Holman & Wiegert developed fitting formulae for both S-type and P-type planets in binary systems given as

$$\frac{a_{\rm cr}}{a} \ = \ 0.464 - 0.38\mu + \mathcal{F}_{\rm S}(\mu, e_{\rm b}) \tag{42}$$

and

$$\frac{a_{\rm cr}}{a} \ = \ 1.60 + 4.12\mu + \mathcal{F}_{\rm P}(\mu, e_{\rm b}) \ , \tag{43}$$

respectively.

These equations give the critical semimajor axis $a_{\rm cr}$ in units of the semimajor axis a in case of S-type and P-type orbits. For an S-type orbit, the ratio $a_{\rm cr}/a$, see Eq. (42), conveys the *upper limit* of planetary orbital stability, whereas for a P-type orbit, the ratio $a_{\rm cr}/a$, see Eq. (43), conveys the *lower limit* of planetary orbital stability. Moreover, μ denotes the stellar mass ratio given as $\mu = M_2/(M_1{+}M_2)$, where M_1 and M_2 constitute the two masses of the binary components with $M_2 \leq M_1$. Equations (42) and (43) also contain the parameter functions $\mathcal{F}_{\rm S}(\mu, e_{\rm b})$ and $\mathcal{F}_{\rm P}(\mu, e_{\rm b})$, which depend on the aforementioned mass ratio μ and the eccentricity of the stellar binary, $e_{\rm b}$. Considering that this paper is solely aimed at stellar binaries in circular orbits (i.e., $e_{\rm b} = 0$), it is found that $\mathcal{F}_{\rm S} = \mathcal{F}_{\rm P} = 0$.

Planetary orbital stability has been investigated by many authors using chaos indicators, such as the maximal Lyapunov exponents (MLE), fast Lyapunov indicator (FLI), and the mean exponential growth factor of nearby orbits (MEGNO), to name those commonly used; see, e.g., Satyal et al. (2013) for details, recent applications, and references. These methods have also been used to characterize the transition from stable to unstable orbits within the framework of the circular and elliptical 3-body problems; see, e.g., Cuntz et al. (2007), Eberle et al. (2008), and Szenkovits & Makó (2008) for details.

In our previous work (e.g., Musielak et al. 2005), we studied the stability of both S-type and P-type orbits in stellar binary systems, and deduced orbital stability limits for planets. These limits were found to depend on the mass ratio between the stellar components and the distance ratio between planetary and binary semimajor axes. This topic was revisited by Eberle et al. (2008), who used the concept of Jacobi's integral and Jacobi's constant to deduce stringent criteria for the stability of planetary orbits in binary systems for the special case of the coplanar circular restricted three-body problem. Recently we studied planetary orbital stability through the perspective of a chaos indicator, the MLE (Quarles et al. 2011). From the use of a chaos indicator a cutoff value for the maximum Lyapunov exponent was determined as an additional stability criterion for S-type planets in the circular restricted 3-body problem.

5. Applications and Classifications

5.1. Case Studies of S-Type Habitability in Binaries

In the following, we provide selected case studies of S-type habitability for binary systems, including systems of equal and non-equal masses (see Table 4). Our intent is to demonstrate the functionality of the method-as-proposed[5]; there is no intent to offer detailed parameter studies of systems or give comparisons to other work; these efforts will be pursued in later papers of this series.

Figure 4 allows comparative insight into S-type habitability for binary systems. The selected cases are systems with masses of $M_1 = M_2 = 1.0\ M_\odot$ and $M_1 = 1.5\ M_\odot$, $M_2 = 1.0\ M_\odot$ with binary separation distances of 10 AU and 20 AU. For single stars of 1.0 $M_\odot$, the CHZ extends from 1.049 to 1.498 AU, and the GHZ extends from 0.927 to 1.831 AU; these values are modestly higher than those for G2 V stars given in Table 3 owing to a minuscule difference in mass (i.e., 0.99 versus 1.0 $\mathrm{M}_\odot$). For an equal-mass binary system of 1.0 $M_\odot$ with a separation distance ($2a$) of 10 AU, the radiative CHZ and GHZ extend from 1.056 to 1.511 AU, and from 0.932 to 1.853 AU, respectively, for each component; the extent of the radiative EHZ is altered from 2.64 to 2.70 AU. However, there is now an (upper) orbital stability limit of 1.37 AU imposed on each star; consequently, significant portions of the radiative CHZ and GHZ are unavailable for providing habitability. If the second star is placed at a distance of 20 AU, the changes in extent for the radiative CHZ and GHZ relative to the single star case are extremely minor. Specifically, for binary separations of 10 AU and 20 AU, the sizes of the radiative GHZ increased by 1.9% and 0.6% relative to the case of a single star case. Moreover, for the system with a separation distance of 20 AU, the imposed orbital stability limit is at 2.74 AU; consequently, the full extents of the radiative CHZ, GHZ, and EHZ are now available for planetary habitability.

---

[5] The method has also successfully been used for determining the radiative habitable zone of Kepler-16, a binary system with $M_1 = 0.69\ M_\odot$ and $M_2 = 0.20\ M_\odot$, and a Saturnian planet in a P-type orbit; see Quarles et al. (2012) for a detailed study of the system's habitability.

Figure 4 also shows results for the pairs $M_1 = 1.5\ M_\odot$ and $M_2 = 1.0\ M_\odot$. In case of a single 1.5 $M_\odot$ mass star, the CHZ and GHZ extend between 1.88 and 2.49 AU, and between 1.65 and 3.11 AU, respectively, whereas the EHZ extends up to 4.61 AU. In this type of system, a secondary star of 1.0 $M_\odot$ placed at a separation distance of 10 AU again modifies the extents of the radiative CHZ and the radiative GHZ, which are now given as from 1.89 to 2.49 AU and from 1.66 to 3.14 AU, respectively; however, there is an imposed orbital stability limit at 1.56 AU. Therefore the entire extents of the radiative CHZ, GHZ, and EHZ of the primary are unavailable to the facilitation of planetary habitability. If the secondary star is placed at a separation distance of 20 AU, the radiative CHZ, GHZ, and EHZ of the primary star are again similar to those of a 1.5 $M_\odot$ mass star. However, the orbital stability limit now occurs at a distance of 3.12 AU from the primary; therefore, the entire radiative EHZ (if measured from the outer edge of the GHZ) is available for habitability.

In summary, for potentially habitable S-type binaries, due to the nature of the relatively large separation of the stellar components, the effect of the stellar secondary on the extents of the RHZs is usually very minor, i.e., about a few percent or less, with the biggest impact occurring in F-type systems. The secondary's main influence on circumstellar habitability consists in imposing restrictions on the facilitation of habitability due to its gravitational influence, which is most relevant in close binary systems; see Sect. 5.3 for additional results. The results of the present work refer to the special case of binaries in circular orbits. More detailed results, which also encompass binary systems in elliptical orbits, thus entailing much more intricate outcomes, have been given by, e.g., Eggl et al. (2012); these types of studies will be revisited in Paper II.

5.2. Case Studies of P-Type Habitability in Binaries

Various sets of models have also been pursued for P-type habitability (see Fig. 6). In this case, we focus on systems with masses of $M_1 = M_2 = 1.0\ M_\odot$ and $M_1 = 1.5\ M_\odot$ and $M_2 = 0.5\ M_\odot$; additionally, we also considered $M_1 = 1.25\ M_\odot$ and $M_2 = 0.75\ M_\odot$ (see Tables 4, 5, 6, and 7 for details). The separation distances ($2a$) were chosen as 0.5, 1.0, and 2.0 AU. Our approach is again a two-step process. We first explore the existence and extent of the radiative CHZ, GHZ, and EHZ; subsequently, we consider the additional constraint of planetary orbital stability, which in the case of P-type orbits constitutes a lower limit (see Sect. 4). The results for the occurrence of habitability can be summarized as follows.

For systems with masses of $M_1 = M_2 = 1.0\ M_\odot$, the following behavior occurs: For separation distances ($2a$) of 0.5 AU, the inner limit (i.e., RHL; see Sect. 3.4) of the radiative CHZ ($s_\ell = 0.95$ AU) is found to vary between 1.46 and 1.54 AU, with 1.54 AU being the acceptable inner limit value; see Eq. (32). Furthermore, the outer limit of the radiative CHZ ($s_\ell = 1.37$ AU) varies between 2.10 and 2.16 AU, where 2.10 AU constitutes the acceptable limit value. In consideration of the orbital stability limit at 0.915 AU, the entire extent of the radiative CHZ offers circumbinary habitability. The acceptable inner limit of the radiative GHZ is given at 1.38 AU, whereas the acceptable outer limit occurs at 2.58 AU; thus, also the entire radiative GHZ is identified as habitable.

For separation distances of 1.0 AU, the orbital stability limit is given at 1.83 AU, which falls inside the domain of the radiative CHZ ranging from 1.69 to 2.06 AU; therefore, only half of the radiative CHZ is available for circumbinary habitability, whereas the other half is not. Noting that the radiative CHZ is fully embedded into the radiative GHZ, entails that also only a fraction of the radiative GHZ is available for circumbinary habitability. However, the full extent of the radiative EHZ, which has an inner limit at 3.70 AU and an outer limit at 3.83 AU is able to offer habitability, if existing. We also studied models with binary separations of 2.0 AU. In this case, the orbital stability limit is found at 3.66 AU. Therefore, both the radiative CHZ and GHZ are unavailable for providing habitability; the latter has an outer limit that varies between 2.39 and 3.05 AU with 2.39 AU constituting the appropriate value. The radiative EHZ is found to vary between 3.60 and 4.09 AU. With 3.60 AU to be ruled acceptable as the conservatively selected (i.e., inner) limit of the EHZ, none of the radiative EHZ should be considered acceptable for circumbinary habitability.

Further case studies were pursued for systems with unequal distributions of masses, and by implication, luminosities as, e.g., systems with $M_1 = 1.5\ M_\odot$ and $M_2 = 0.5\ M_\odot$. Following the well-known mass–luminosity relationship for main-sequence stars, it is found that a 1.5 $M_\odot$ star possess a luminosity L_* about 3.5 times higher than a 1.0 $M_\odot$ star (see Table 4); a similar factor of difference applies to the recast stellar luminosity L' (see Sect. 3.1). Thus, the combined luminosity of the $(1.5\ M_\odot, 0.5\ M_\odot)$ system is considerably higher than the combined luminosity of the $(1.0\ M_\odot, 1.0\ M_\odot)$, as expected, and the distribution of luminosities is highly uneven. On the other hand, following the formula of Holman & Wiegert (1999), as well as previous work by Dvorak (1986), an unequal distribution of stellar mass, i.e., a smaller value of μ (see Sect. 4), entails a smaller orbital stability limit. Because it constitutes a lower limit, i.e., positioned relatively close to the two stars, it offers a relatively large "window of opportunity" for planets in any of the RHZs to be orbitally stable.

Results are given in Fig. 6. For a binary separation of 0.5 AU it is found that both the radiative CHZ and GHZ exist, and they are both fully permitted based on the planetary orbital stability constraint, although the width of the CHZ is relatively small. The CHZ extents from 2.14 to 2.28 AU, whereas the GHZ extents from 1.91 to 2.90 AU; the orbital stability limit is given at 0.66 AU. There is also a considerably large domain of the EHZ, which has an inner limit at 4.40 AU and an outer limit at 4.89 AU.

However, for binary separation of 1.0 AU, the radiative CHZ has been nullified, even though its existence would be consistent with the planetary orbital stability constraint conveying a lower limit at 1.32 AU. The reason is that the RHL of s_4, which is at 2.38 AU, is located below the RHL of s_2, given as 2.05 AU. The same criterion (see Eq. 36) also dictates the relatively small width of the radiative GHZ, which extends between 2.16 and 2.66 AU. At a binary separation of 2.0 AU, the situation is even more drastic as both the radiative CHZ and GHZ are disallowed; the only type of circumbinary habitability remaining is that provided by the relatively large EHZ, which is also consistent with the planetary orbital stability constraint.

Additional analyses have been pursued with the binary separation distance $2a$ taken as an independent variable. Hence, for systems of $M_1 = M_2 = 1.0\ M_\odot$, it is found that the radiative CHZ and GHZ vanish at $2a = 1.52$ AU and 2.35 AU, respectively; the associated limits of orbital stability are given as 2.78 and 4.31 AU, respectively. Furthermore, the distances of fading away for the radiative CHZ and GHZ are identified as 1.76 and 2.45 AU, respectively. Therefore, it can be concluded that for equal-mass binary systems with $M_1 = M_2 = 1.0\ M_\odot$, habitability for relatively wide binaries is lost because of the lack of planetary orbital stability occurring at binary separations with the circumbinary RHZ still present.

We repeated the same type of study for binary systems with $M_1 = 1.5\ M_\odot$ and $M_2 = 0.5\ M_\odot$. In this case it was found that the radiative CHZ and GHZ vanish at $2a = 0.54$ AU and 1.24 AU, respectively; here the associated limits of orbital stability are given at 0.71 and 1.63 AU, respectively. Also, the distances of fading away regarding the radiative CHZ and GHZ are found at 1.28 and 2.35 AU, respectively. For this type of system it is found that habitability for relatively wide binaries is lost because of the lack of existence of the circumbinary RHZ, even though the orbital stability criterion would still be met. The fact that the circumbinary RHZ starts to fade away at relatively small binary distances, if compared to the case $M_1 = M_2 = 1.0\ M_\odot$, is relatively surprising in the view of the overall much higher system luminosity because of M_1 and $L_* \propto M_*^4$ (e.g., Reid 1987); see Sect. 2. However, it is a consequence of the imbalance of binary systems with highly unequal distributions of stellar luminosity outputs.

Radiative imbalance of binary systems in the context of circumbinary habitability is strongly related to a property previous discussed, which is that it may occur that the RHL for $s_{\ell,\mathrm{out}}$ to be partially or completely located inside of the RHL for $s_{\ell,\mathrm{in}}$, see Sect. 3.4. Therefore, we also report more detailed results about the variation of the inner and outer limits (RHLs), which if appropriate amount to the establishment of RHZs; see Eq. (32). Specifically, as conveyed through Eq. (36) for a specific RHZ to exist, it is required for the

RHL of $s_{\ell,\mathrm{out}}$ to be located completely outside of the RHL for $s_{\ell,\mathrm{in}}$ with the $s_{\ell,\mathrm{in}}$ and the $s_{\ell,\mathrm{out}}$ being appropriately paired.

The condition of Eq. (36) is most readily met in case of equal-mass binary systems with small separation distances; e.g., complete separation for the inner and outer limits for the appropriate ($s_{\ell,\mathrm{in}}, s_{\ell,\mathrm{out}}$ pairings is found for, e.g., $M_1 = M_2 = 1.0\ M_\odot$ and $a = 0.25$ AU. Conversely, the nullification of RHZs is most likely to occur in wide binary systems with unequal masses. Figure 7 depicts the ranges of the various RHLs for various cases previously discussed. It shows that for $M_1 = M_2 = 1.0\ M_\odot$ and $a = 0.25$ AU. the RHLs pertaining to the radiative CHZ and GHZ are wiedely separated; thus, the radiative CHZ and GHZ are clearly able to exist.

For $M_1 = 1.5\ M_\odot$, $M_2 = 0.5\ M_\odot$ and $a = 0.25$ AU the pairings (s_2, s_4) and (s_1, s_5) are again separated, though for (s_2, s_4) this is barely the case as the upper limit of the s_2 range almost coincides with the lower limit of the s_4 range; nevertheless, the full extents of the radiative CHZ and GHZ are permitted, and possibility of habitability for these types of RHZs solely depends on the fulfillment of the planetary orbital stability constraint. However, in case of $M_1 = 1.5\ M_\odot$, $M_2 = 0.5\ M_\odot$ and $a = 1.0$ AU both the radiative CHZ and GHZ are disallowed. In this case, only the radiative EHZ is permitted. Fortunately, it is also permitted based on the planetary orbital stability constraint (see Fig. 6), implying that this type of system offers a tenuous chance of circumbinary habitability.

Although the unequal distribution of stellar mass is identified as highly advantageous to the facilitation of planetary orbital stability, in consideration of the lower stability limit for P-type orbits for smaller mass ratios μ (see Sect. 4), the situation for the existence of the RHZs is much less ideal, even for systems where the stellar primary is highly luminous. In this regard the radiative CHZ is in most jeopardy as $(s_4 - s_2)$ constitutes the smallest bracket among the various kinds of HZs (see Table 2). More fortunate situations are found for the radiative GHZ and EHZ, which are also characterized by larger widths, especially in systems of stars with high luminosities.

5.3. Habitability Classification: Habitability Types S, P, ST, and PT

Another key aspect of this study is provide an appropriate classification of circumstellar habitability, which will be illustrated taking equal-mass binary systems for theoretical main-sequence stars of spectral types F0 to M0 as examples. The required stellar parameters, which are the effective temperatures T_{eff}, the luminosities L_*, and the masses M_* of the stellar components are given in Table 1.

Previously, Dvorak (1982) introduced the terminology of S-type and P-type orbits for system planets, which is now widely used by the orbital stability, planetary, and the astrobiology science communities. Evidently, besides the assessment of orbital stability behaviors, these terms are also appropriate for classifying binary system RHZs, if existing. However, following previous investigations (e.g., Takeda et al. 2008; Dvorak et al. 2010; Haghighipour et al. 2010; Kopparapu & Barnes 2010; Eggl et al. 2012), as well as the results of the present work, the spatial domain of S-type and P-type habitability depicted by the RHZSs is often adversely affected, and in some cases even nullified, by the requirement that system planets must be orbitally stable. Thus, if the available extent of the S-type and P-type RHZSs for the manifestation of habitability is truncated owing to the additional constraint of planetary orbital stability, these zones will be referred to as ST-type and PT-type, respectively, in the following.

Computationally, the occurrence of P, PT, ST, and S-type habitability can be identified as follows (see Fig. 8). For sufficiently small binary separations ($2a$), it is found that both the inner and outer limit of the P-type RHZ are located outside of the P-type orbital stability limit (see Eq. 43), which constitutes a *lower limit* of planetary orbital stability (see Sect. 4). If the distance of binary separation is increased, the inner and outer limits of the P-type RHZ decrease, whereas the P-type orbital stability limit increases. Thus, starting at a certain value of $2a$, the inner limit of the P-type RHZ will be positioned inside of the P-type orbital stability limit; in this case, PT-type habitability is attained. If the binary separation is further increased, both the inner and outer limit of the P-type RHZ (if existing) will move inside the P-type orbital stability limit; thus, there will be no habitability.

For mid-sized values of the binary separation distance, the inner and outer limit of the S-type RHZ are located inside of the S-type orbital stability limit (see Eq. 42), which constitutes an *upper limit* of planetary orbital stability (see Sect. 4). This setting again entails the absence of habitability for binary system planets. If the binary separation distance is further increased, the S-type orbital stability will increase and will move beyond the inner limit of the S-type RHZ; in this case, ST-type habitability is encountered. Eventually, for sufficiently large binary separations, the S-type orbital stability limit is positioned at a sufficiently large distance from the stellar component in order for both the inner and outer limit of the S-type RHZ, which moved closer to the stellar component, to be located inside the S-type orbital stability limit. Now, S-type habitability is found because the total width of the S-type RHZ is available for facilitating habitability.

S-type, P-type, ST-type, and PT-type habitability for the CHZ, GHZ, and EHZ for equal-mass main-sequence star binaries of spectral types F0 to M0 are displayed in Tables 8, 9, and 10. The CHZ, GHZ, and EHZ correspond to the habitability limits (s_2, s_4), (s_1, s_5), and (s_1, s_6), respectively, see Sect. 3.4. It is found that for the evaluated array of spectral type / binary separation combinations, with the latter inspected between distances $(2a)$ of 0.2 and 10.0 AU, the CHZ exhibits the most cases of non-existent habitability, as expected, owing to its property that it has the smallest extent compared to the GHZ and EHZ. But if habitability for the CHZ is encountered, it is usually of type S or P rather than ST or PT, noting that because of the CHZ's small extent, cutting through (i.e., truncation) of the RHZ by the S-type or P-type orbital stability limit (see Eqs. 42 and 43) is relatively rare.

On the other hand, for the GHZ and the EHZ, owing to their significantly larger extents, there is a much smaller range of binary separation distances without the possibility of circumstellar habitability. However, if habitability is realized, there is a higher percentage of PT versus P cases, as well as ST versus S cases, found for the GHZ, and even more so for the EHZ, compared to the CHZ. This behavior is again due to the fact that the GHZ, and even more so the EHZ, is much more extended than the EHZ. Thus, there is a higher likelihood that the S-type or P-type planetary orbital stability limit interferes with those zones. Most importantly, there is also no case of an equal-star binary system between the spectral types F0 and M0 where both S/ST or P/PT-type habitability is found to be possible. This result has already been suggested by the Eqs. (14) and (16), see Sect. 3.2.

For G2 V stars, i.e., two Suns in a circular orbit, the following picture emerges: For the CHZ, P-type habitability is encountered up to a distance of 0.6 AU, followed by PT-type habitability up to a distance $(2a)$ of 0.8 AU. ST-type habitability is encountered at a binary separation of 8.3 AU, whereas S-type habitability occurs at 12.0 AU. Regarding the GHZ and EHZ, somewhat smaller values compared to the CHZ occur for P-type habitability, and moderately larger values occur for PT-type habitability. The biggest difference between the GHZ and EHZ, on one hand, and the CHZ, on the other hand, is found for S-type habitability: For the GHZ and EHZ, this mode of habitability requires binary separations of 14.6 and 21.0 AU, respectively; see Table 11 for complete results.

Another representation of our results is given in Fig. 9. It conveys a depiction of S/ST-type and P/PT-type habitability using different gray scales regarding the CHZ, GHZ, and EHZ. It also shows a notable domain with no habitability at all. Note that no separate S/ST-type contour is attained for the EHZ, owing to the fact that the inner boundary of the EHZ (i.e., s_1; see Table 2) agrees with the inner boundary of the GHZ, thus rendering the same criterion for the inner limit of habitability. For the distance limits (2a) of ST-type habitability at the high-luminosity end of the stellar range (attained outside the figure's frame), the values for the F5 V, F2 V, and F0 V stars pertaining to the GHZ and EHZ are given as 11.7, 13.6, and 15.9 AU, whereas for the CHZ, they are given as 13.3, 15.5, and 18.1 AU, respectively.

6. Summary and Conclusions

This study is aimed at developing a new method for a comprehensive assessment of S-type and P-type habitability in stellar binary systems. P-type orbits occur when the planet orbits both binary components, whereas in case of S-type orbits the planet orbits only one of the binary components with the second component considered a perturbator. An important characteristic of the method is that the joint constraint of orbital stability for a system planet as well as the necessity for providing a habitable environment for that planet through the stellar radiative energy fluxes ("radiative habitable zone") needs to be met, a property also recognized in previous studies (e.g., Takeda et al. 2008; Kopparapu & Barnes 2010; Eggl et al. 2012; Kane & Hinkel 2013).

In conjunction with the present study, a habitability classification of type S, P, ST, and PT is also proposed, which considers whether or not S-type and P-type radiative habitable zones are truncated based on the additional constraint of planetary orbital stability. This classification scheme may be of general relevance as it can be applied to both equal-mass and non-equal mass binary systems, as well as to systems with binaries in elliptical orbits.

A key aspect of the proposed method is the introduction of a combined algebraic formalism for the assessment of both S-type and P-type habitability; in particular, mathematical criteria are presented allowing to determine for which systems S-type and P-type RHZs are realized. In this regard, apriori choices about the presence of S-type and P-type RHZs are neither necessary nor possible as the existence of S-type as well as P-type RHZs is proliferated through well-defined mathematical conditions regarding the underlying descriptive fourth-order algebraic equation. The coefficients of the polynomial are given by the binary separation distance ($2a$), the solar system-based parameter for the limit of habitability (s_ℓ), and the modified values for the luminosities (L'_1, L'_2) of the stellar binary components. Regarding the binary system habitable zone, we consider conservative, general and extended zones of habitability, noting that their inner and outer limits are informed by previous solar system investigations (e.g., Kasting et al. 1993; Underwood et al. 2003; Selsis et al. 2007).

After combining the requirements pertaining to the existence of the RHZs with the necessity of orbital stability for system planets, we were able to identify five different cases, which are: S-type and P-type habitability provided by the full extent of the RHZ; habitability, where the RHZ is truncated by the additional constraint of planetary orbital stability (labelled as ST and PT-type, respectively); and cases of no habitability at all. This will allow a general classification of habitability which is expected to be also useful for higher-order systems, i.e., $N > 2$. We provided examples for various cases, i.e., equal-mass and nonequal-mass systems, by focusing on theoretical main-sequence stars based on parameters given by Castelli & Kurucz (2004), Gray (2005), and Kurucz (2005). As a limitation of scope, we restricted our investigation to binary systems with the stellar components in circular orbits; note that systems in elliptical orbits will be examined in Paper II.

In the segment of applications, we examined the existence of habitable S-type orbits for selected examples. We found that regarding the RHZs, owing to the relatively large separation of the stellar components, the effect of the stellar secondary on the extents of the RHZs is typically very minor. The secondary's main influence on circumstellar habitability consists in imposing restrictions regarding planetary orbital stability in form of an upper stability limit around each stellar component, which often truncates or nullifies S-type planetary habitability. Previous studies, which also include binary systems in elliptical orbits, thus entailing much more intricate outcomes, were given by, e.g., Eggl et al. (2012). In the framework of our study, we specifically considered the radiative EHZ. It was found that this kind of zone is most affected by the limitation of planetary orbital stability as it is closest to the secondary stellar component.

More complicated scenarios emerge for potentially habitable P-type orbits, which were studied in detail through comparing pairs with masses of (1.0 $M_\odot$, 1.0 $M_\odot$) to pairs with masses of (1.5 $M_\odot$, 0.5 $M_\odot$). Results for pairs of (1.25 $M_\odot$, 0.75 $M_\odot$) were given as intermediate case for additional information. The overall focus of this effort was to deduce information on how the distribution of stellar mass and, by implication, stellar luminosity impacts the habitability prospects of main-sequence star binary systems. It was found that although the unequal distribution of stellar mass is highly advantageous to the facilitation of planetary orbital stability, in consideration of the lower stability limit for P-type orbits (see Sect. 4), the situation for the existence of the RHZs may be highly complicated.

The reason is that an unequal distribution of mass always entails un unequal distribution of luminosity with the more massive stellar component being overwhelmingly luminous in accord to the mass–luminosity relationship, i.e., $L_* \propto M_*^4$ (Reid 1987); see Sect. 2. Generally, this property drastically increases the overall habitability prospects of a P-type binary system. However, there is an important complication that is particularly relevant for relatively wide binaries: Careful evaluations show that for some systems the outer limit of distinct RHZs may be positioned inside the inner limit of that zone, at least in some parts of the system (see, e.g., Fig. 8) , which effectively nullifies that zone, irrespectively of planetary orbital stability considerations. In this regard, the CHZ is in highest jeopardy as $(s_4 - s_2)$ constitutes the smallest bracket among the various kinds of HZs (see Table 2).

Future work will deal with a significant augmentation of our method to other systems, including systems with binary components in elliptical orbits (see Paper II). This will allow us to compare the potential and outcome of our method, as well as results for individual systems, to other findings in the literature. We also expect our method to be applicable to general binary systems with main-sequence stars as well as to systems containing evolved stars; this latter effort is motivated by observational evidence and supporting theoretical efforts indicating that planets are also able to exist around stars that have left the main-sequence (e.g., Sato et al. 2003; Ramm et al. 2009; Eberle & Cuntz 2010; Doyle et al. 2011). Particularly, it is highly desirable to augment our method to systems of higher order, as motivated by the steady progress in theory as well as ongoing and future observational discoveries of exosolar planetary systems.

This work has been supported in part by NASA through the American Astronomical Society's Small Research Grant Program. The author acknowledges comments by B. Quarles and Z. E. Musielak as well as assistance with computer graphics by S. Sato and S. Satyal. This study made use of the software applications Fortran®, MATHEMATICA®, and MATLAB®. The author anticipates the development of a black box code, called `BinHab`, to be hosted at The University of Texas at Arlington, which will allow the assessment of habitability in binary systems based on the developed method.

REFERENCES

Abramowitz, M., & Stegun, I. A. 1972, Handbook of Mathematical Functions with Formulas, Graphs, and Mathematical Tables, 9th printing (New York: Dover)

Asghari, N., Broeg, C., Carone, L., et al., 2004, A&A, 426, 353

Beyer, W. H. 1987, Handbook of Mathematical Sciences, 6th edn. (Boca Raton: CRC Press)

Bonavita, M., & Desidera, S. 2007, A&A, 468, 721

Bronshtein, I. N., & Semendyayev, K. A. 1997, Handbook of Mathematics, 3rd edn. (New York: Springer)

Castelli, F., & Kurucz, R. L. 2004, in IAU Symp. 210, Modelling of Stellar Atmospheres ed. N. E. Piskunov, W. W. Weiss, & D. F. Gray (San Francisco: ASP), CD-ROM Poster 20; arXiv:astro-ph/0405087

Cuntz, M., & Yeager, K. E. 2009, ApJ, 697, L86

Cuntz, M., Eberle, J., & Musielak, Z. E. 2007, ApJ, 669, L105

Doyle, L. R., Carter, J. A., Fabrycky, D. C., et al. 2011, Science, 333, 1602

Duquennoy, A., & Mayor, M. 1991, A&A, 248, 485

Dvorak, R. 1982, Österreichische Akademie der Wissenschaften, Mathematisch-Naturwissenschaftliche Klasse, Sitzungsberichte Abt. 2, 191 (10), 423

Dvorak, R. 1984, Celest. Mech. Dyn. Astron., 34, 369

Dvorak, R. 1986, A&A, 167, 379

Dvorak, R., Pilat-Lohinger, E., Bois, E., et al. 2010, Astrobiology, 10, 33

Eberle, J., & Cuntz, M. 2010, ApJ, 721, L168

Eberle, J., Cuntz, M., & Musielak, Z. E. 2008, A&A, 489, 1329

Eggenberger, A., Udry, S., & Mayor, M. 2004, A&A, 417, 353

Eggenberger, A., Udry, S., Chauvin, G., Beuzit, J.-L., Lagrange, A.-M., Ségransan, D., & Mayor, M. 2007, A&A, 474, 273

Eggl, S., Pilat-Lohinger, E., Georgakarakos, N., Gyergyovits, M., & Funk, B. 2012, ApJ, 752, A74

Fatuzzo, M., Adams, F. C., Gauvin, R., & Proszkow, E. M. 2006, PASP, 118, 1510

Forget, F., & Pierrehumbert, R. T. 1997, Science, 278, 1273

Gray, D. F. 2005, The Observation and Analysis of Stellar Photospheres, 2nd edn. (Cambridge: Cambridge University Press)

Haghighipour, N., Dvorak, R., & Pilat-Lohinger, E. 2010, in Planets in Binary Systems, ed. N. Haghighipour, Astrophysics and Space Science Library, Vol. 366 (New York: Springer Science + Business Media), p. 285

Halevy, I., Pierrehumbert, R. T., & Schrag, D. P. 2009, J. Geophys. Res., 114, D18112

Holman, M. J., & Wiegert, P. A. 1999, AJ, 117, 621

Jackson, J. D. 1999, Classical Electrodynamics, 3rd edn., John Wiley & Sons, Inc., Hoboken, NJ

Jones, B. W., Sleep, P. N., & Chambers, J. E. 2001, A&A, 366, 254

Jones, B. W., & Sleep, P. N. 2010, MNRAS, 407, 1259

Kane, S. R., & Hinkel, N. R. 2013, ApJ, 762, A7

Kasting, J. F., Whitmire, D. P., & Reynolds, R. T. 1993, Icarus, 101, 108

Kopparapu, R. K., & Barnes, R. 2010, ApJ, 716, 1336

Kurucz, R. L. 2005, Mem. S. A. It., 8, 14

Lada, C. J. 2006, ApJ, 640, L63

Menou, K., & Tabachnik, S. 2003, ApJ, 583, 473

Mischna, M. A., Kasting, J. F., Pavlov, A., & Freedman, R. 2000, Icarus, 145, 546

Musielak, Z. E., Cuntz, M., Marshall, E. A., & Stuit, T. D. 2005, A&A, 434, 355

Noble, M., Musielak, Z. E., & Cuntz, M. 2002, ApJ, 572, 1024

Orosz, J. A., Welsh, W. F., Carter, J. A., et al. 2012, Science, 337, 1511

Patience, J., White, R. J., Ghez, A. M., et al. 2002, ApJ, 581, 654

Perryman, M. 2011, The Exoplanet Handbook (Cambridge: Cambridge University Press)

Quarles, B., Eberle, J., Musielak, Z. E., & Cuntz, M. 2011, A&A, 533, A2

Quarles, B., Musielak, Z. E., & Cuntz, M. 2012, ApJ, 750, A14

Quintana, E. V., & Lissauer, J. J. 2010, in Planets in Binary Systems, ed. N. Haghighipour, Astrophysics and Space Science Library, Vol. 366 (New York: Springer Science + Business Media), p. 265

Raghavan, D., Henry, T. J., Mason, B. D., et al. 2006, ApJ, 646, 523

Raghavan, D., McAlister, H. A., Henry, T. J., et al. 2010, ApJS, 190, 1

Ramm, D. J., Pourbaix, D., Hearnshaw, J. B., & Komonjinda, S., 2009, MNRAS, 394, 1695

Reid, N. 1987, MNRAS, 225, 873

Sándor, Zs., Sűli, Á., Érdi, B., Pilat-Lohinger, E., Dvorak, R. 2007, MNRAS, 375, 1495

Sato, B., et al., 2003, ApJ, 597, L157

Satyal, S., Quarles, B., & Hinse, T. C. 2013, MNRAS, submitted

Selsis, F., Kasting, J. F., Levrard, B., Paillet, J., Ribas, I., & Delfosse, X. 2007, A&A, 476, 1373

Setiawan, J., Hatzes, A. P., von der Lühe, et al. 2003, A&A, 398, L19

Slawson, R. W., Prša, A., Welsh, W. F., et al. 2011, AJ, 142, 160

Stix, M. 2004, The Sun: An Introduction, 2nd edn. (A&A Library, Berlin: Springer)

Szenkovits, F., & Makó, Z. 2008, Celest. Mech. Dyn. Astron., 101, 273

Takeda, G., Kita, R., & Rasio, F. A. 2008, ApJ, 683, 1063

Trilling, D. E., Stansberry, J. A., Stapelfeldt, K. R., et al. 2007, ApJ, 658, 1289

Underwood, D. R., Jones, B. W., & Sleep, P. N. 2003, Int. J. Astrobiol., 2, 289

Williams, D. M., & Pollard, D. 2002, Int. J. Astrobiol., 1, 61

This preprint was prepared with the AAS LaTeX macros v5.0.



Fig. 1.— Set-up for the mathematical treatment of S-type (top) and P-type (bottom) habitable zones of binary systems as given by the stellar radiative fluxes. Note that the stars S1 and S2 have been depicted as identical for convenience.



Fig. 2.— Habitability limits for equal-mass binaries regarding S-type and P-type planetary orbits pertaining to the RHZ. The various limits, separated by grayish areas, refer to different values of s_ℓ given as 0.84, 0.95, 1.00, 1.37, 1.67, and 2.40 AU (from left to right; corresponding to $\ell = 1$ to 6); see Sect. 2 and 3.2 for details.



Fig. 3.— Examples of S-type and P-type RHZs for different systems obtained through solutions of the governing fourth-order polynomial equations. The thick solid lines indicate the RHLs, corresponding to the inner and outer limit of habitability based on $s_\ell = 0.84$ and 1.67 AU, respectively. Note that in the S-type case the RHLs are bended toward the center, whereas in the P-type case they are of elliptical shape. The top row displays systems with $L_1' = L_2' = L_\odot$, whereas the bottom top row displays systems with $L_1' = 1.5\ L_\odot$ and $L_2' = 0.5\ L_\odot$. The left column assumes separation distances ($2a$) of 5.0 AU rendering P-type RHZs, whereas the right column assumes separation distances of 0.5 AU resultant in S-type RHZs. The gray areas indicate the appropriate circular regions (annuli), referred to as RHZs, for each case.



Fig. 4.— Maximum of permissible separation distance $2a$ of equal-mass binary components, which still permits the possibility of P-type RHZs (see Eq. 41). Results are given for the CHZ, GHZ, and EHZ.



Fig. 5.— Domains of S-type habitability for main-sequence star binary systems. The left column depicts systems with masses of $M_1 = M_2 = 1.0\ M_\odot$, whereas the right column depicts systems with masses of $M_1 = 1.5\ M_\odot$ and $M_2 = 1.0\ M_\odot$. The separation distances ($2a$) are given as 10 AU (top) and 20 AU (bottom). Results for the conservative, general and extended RHZs are given as dark gray, medium gray, and light gray areas, respectively. The orbital stability limits are indicated by red dashed lines.



Fig. 6.— Domains of P-type habitability for main-sequence star binary systems. The left column depicts systems with masses of $M_1 = M_2 = 1.0\ M_\odot$, whereas the right column depicts systems with masses of $M_1 = 1.5\ M_\odot$ and $M_2 = 0.5\ M_\odot$. The separation distances ($2a$) are given as 0.5 AU (top), 1.0 AU (middle), and 2.0 AU (bottom). Results for the conservative, general and extended RHZs are given as dark gray, medium gray, and light gray areas, respectively. The orbital stability limits are indicated by red dashed lines.



Fig. 7.— Ranges of the inner and outer limits of RHZs, given as $\mathrm{Min}\Big(\mathcal{R}(z,\varphi)\Big)\Big|_{s_\ell}$ and $\mathrm{Max}\Big(\mathcal{R}(z,\varphi)\Big)\Big|_{s_\ell}$, pertaining to $s_\ell = 0.84$, 0.95, 1.37, 1.67, and 2.40 AU (from left to right; see labels 1, 2, 4, 5, and 6). The figure panel depicts binary systems with separation distances ($2a$) of 0.5 AU (top) and 2.0 AU (bottom) and with masses of $M_1 = M_2 = 1.0\ M_\odot$ (left) and $M_1 = 1.5\ M_\odot$ and $M_2 = 0.5\ M_\odot$ (right); see also Tables 5 to 7 for additional information.



Fig. 8.— Facilitation of habitability in an equal-star binary system of two G2 V stars as function of the binary separation distance ($2a$) pertaining to the GHZ, i.e., $s_{\ell,\mathrm{in}} = 0.84$ and $s_{\ell,\mathrm{out}} = 1.67$ AU. The two red lines indicate the limits of the P-type RHZ (i.e., $\mathrm{RHZ}_{\mathrm{in}}$ and $\mathrm{RHZ}_{\mathrm{out}}$; see Eqs. (40a) and (40b)), whereas the two blue lines indicate the limits of the S-type RHZ (see Eqs. (37a) and (37b)). The available S-type and P-type RHZs are depicted as grayish areas. Furthermore, the cyan dashed line indicates the P-type orbital stability limit, whereas the violet dashed line indicates the S-type orbital stability limit; see Eqs. (43) and (42), respectively. Note that the P-type orbital stability limit constitutes a lower limit (i.e., planetary orbital stability is attained in the domain to the upper left of that line), whereas the S-type orbital stability limit constitutes an upper limit (i.e., planetary orbital stability is attained below that line). Therefore, the intersections between the P-type RHZs and the P-type orbital stability limit allow to identify the domains of P-type and PT-type habitability given to the left of the intersection points. Also, the intersections between the S-type RHZs and the S-type orbital stability limit allow to identify the domains of S-type and ST-type habitability given to the right of the intersection points. No habitability is found between the intersection points PT and ST. Similar figures hold for other types of binary systems, including systems of non-equal stellar masses.



Fig. 9.— Depiction of S/ST-type and P/PT-type habitability based on the joint constraint of planetary orbital stability and the availability of a habitable environment provided by the stellar radiative energy fluxes. The domains of the conservative, general and extended HZs are depicted as dark gray, medium gray, and light gray areas, respectively. The results refer to the particular case of equal-mass binaries in circular orbits, depicted as main-sequence stars of identical spectral types.

Table 1. Stellar Parameters

Sp. Type	$T_{\rm eff}$	R_*	L_*	$S_{{\rm rel},*}$	M_*
...	(K)	$(R_\odot)$	$(L_\odot)$	...	$(M_\odot)$
F0	7178	1.62	6.255	1.145	1.60
F2	6909	1.48	4.481	1.113	1.52
F5	6528	1.40	3.196	1.072	1.40
F8	6160	1.20	1.862	1.037	1.19
G0	5943	1.12	1.405	1.019	1.05
G2	5811	1.08	1.194	1.009	0.99
G5	5657	0.95	0.830	0.997	0.91
G8	5486	0.91	0.673	0.985	0.84
K0	5282	0.83	0.481	0.971	0.79
K2	5055	0.75	0.330	0.957	0.74
K5	4487	0.64	0.149	0.926	0.67
K8	4006	0.53	0.066	0.905	0.58
M0	3850	0.48	0.045	0.900	0.51

Note. — $S_{{\rm rel},*}$ is calculated using $s_\ell = s_3$.

Table 2. Habitability Limits for Solar Environments

ℓ ...	s_ℓ (AU)	Description ...
1	0.84	Runaway greenhouse effect
2	0.95	Start of water loss
3	1.00	Earth-equivalent position
4	1.37	First CO_2 condensation
5	1.67	Maximum greenhouse effect, no clouds
6	2.40	Maximum greenhouse effect, 100% clouds

Note. — See text for references.

Table 3. Habitable Zones of Single Main-Sequence Stars

Sp. Type	Habitable Zone				
...	HZ-iG (AU)	HZ-iC (AU)	HZ-oC (AU)	HZ-oG (AU)	HZ-oE (AU)
F0	1.978	2.253	2.909	3.659	5.485
F2	1.696	1.929	2.543	3.178	4.723
F5	1.456	1.653	2.243	2.780	4.085
F8	1.128	1.278	1.782	2.192	3.188
G0	0.987	1.118	1.584	1.940	2.805
G2	0.915	1.035	1.480	1.808	2.606
G5	0.766	0.866	1.253	1.527	2.192
G8	0.694	0.784	1.148	1.395	1.994
K0	0.590	0.667	0.990	1.199	1.705
K2	0.492	0.555	0.837	1.010	1.429
K5	0.335	0.378	0.593	0.709	0.990
K8	0.224	0.253	0.410	0.487	0.674
M0	0.187	0.211	0.345	0.409	0.565

Note. — HZ-i and HZ-o indicate the inner and outer limits, respectively, of single main-sequence star habitable zones. The added letters C, G and E are in reference to the conservative, general and extended habitable zones, respectively (see text for details).

Table 4. Target List, Sections 5.1 and 5.2

M_* ($M_\odot$)	$T_{\rm eff}$ (K)	L_* ($L_\odot$)	L' ($L_\odot$)	$\mathrm{HZ}_{\oplus,\mathrm{eqv}}$ (AU)
0.50	3857	0.045	0.050	0.22
0.75	5100	0.357	0.372	0.61
1.00	5833	1.228	1.216	1.10
1.25	6256	2.173	2.078	1.44
1.50	6842	4.233	3.830	1.96

Note. — The position of $\mathrm{HZ}_{\oplus,\mathrm{eqv}}$ corresponds to $s_\ell = s_3$.

Table 5. P-Type Stellar Habitability; $a = 0.25$ AU

HZ	Binary System Data		
s_ℓ (AU)	$\mathrm{Min}\Big(\mathcal{R}(z,\varphi)\Big)\Big\vert_{s_\ell}$ (AU)	$\mathrm{Max}\Big(\mathcal{R}(z,\varphi)\Big)\Big\vert_{s_\ell}$ (AU)	$a_{\rm cr}$ (AU)
Model: $M_1 = 1.0\ M_\odot$, $M_2 = 1.0\ M_\odot$			
0.84	1.29	1.38	0.92
0.95	1.46	1.54	0.92
1.00	1.54	1.62	0.92
1.37	2.10	2.16	0.92
1.67	2.58	2.62	0.92
2.40	3.72	3.76	0.92
Model: $M_1 = 1.25\ M_\odot$, $M_2 = 0.75\ M_\odot$			
0.84	1.19	1.52	0.79
0.95	1.36	1.69	0.79
1.00	1.45	1.79	0.79
1.37	1.96	2.28	0.79
1.67	2.43	2.76	0.79
2.40	3.58	3.91	0.79
Model: $M_1 = 1.5\ M_\odot$, $M_2 = 0.5\ M_\odot$			
0.84	1.42	1.91	0.66
0.95	1.65	2.14	0.66
1.00	1.73	2.22	0.66
1.37	2.28	2.76	0.66
1.67	2.90	3.38	0.66
2.40	4.40	4.89	0.66

Table 6. P-Type Stellar Habitability; $a = 0.50$ AU

HZ	Binary System Data		
s_ℓ (AU)	$\mathrm{Min}\Big(\mathcal{R}(z,\varphi)\Big)\Big\vert_{s_\ell}$ (AU)	$\mathrm{Max}\Big(\mathcal{R}(z,\varphi)\Big)\Big\vert_{s_\ell}$ (AU)	$a_{\rm cr}$ (AU)
Model: $M_1 = 1.0\ M_\odot$, $M_2 = 1.0\ M_\odot$			
0.84	1.21	1.54	1.83
0.95	1.40	1.69	1.83
1.00	1.48	1.76	1.83
1.37	2.06	2.28	1.83
1.67	2.54	2.72	1.83
2.40	3.70	3.83	1.83
Model: $M_1 = 1.25\ M_\odot$, $M_2 = 0.75\ M_\odot$			
0.84	1.11	1.75	1.57
0.95	1.29	1.92	1.57
1.00	1.38	2.02	1.57
1.37	1.90	2.49	1.57
1.67	2.35	2.96	1.57
2.40	3.46	4.12	1.57
Model: $M_1 = 1.5\ M_\odot$, $M_2 = 0.5\ M_\odot$			
0.84	1.21	2.16	1.32
0.95	1.43	2.38	1.32
1.00	1.51	2.47	1.32
1.37	2.05	3.00	1.32
1.67	2.66	3.62	1.32
2.40	4.17	5.13	1.32

Table 7. P-Type Stellar Habitability; $a = 1.0$ AU

HZ	Binary System Data		
s_ℓ (AU)	$\mathrm{Min}\Big(\mathcal{R}(z,\varphi)\Big)\Big\|_{s_\ell}$ (AU)	$\mathrm{Max}\Big(\mathcal{R}(z,\varphi)\Big)\Big\|_{s_\ell}$ (AU)	$a_{\rm cr}$ (AU)
Model: $M_1 = 1.0\ M_\odot$, $M_2 = 1.0\ M_\odot$			
0.84	0.85	1.98	3.66
0.95	1.10	2.11	3.66
1.00	1.20	2.18	3.66
1.37	1.87	2.64	3.66
1.67	2.39	3.05	3.66
2.40	3.60	4.09	3.66
Model: $M_1 = 1.25\ M_\odot$, $M_2 = 0.75\ M_\odot$			
0.84	0.70	2.23	3.15
0.95	0.95	2.40	3.15
1.00	1.07	2.50	3.15
1.37	1.69	2.95	3.15
1.67	2.18	3.42	3.15
2.40	3.32	4.55	3.15
Model: $M_1 = 1.5\ M_\odot$, $M_2 = 0.5\ M_\odot$			
0.84	0.83	2.66	2.63
0.95	1.09	2.88	2.63
1.00	1.18	2.96	2.63
1.37	1.74	3.50	2.63
1.67	2.28	4.12	2.63
2.40	3.71	5.63	2.63

Table 8. Assessment of the Conservative Habitable Zone of Equal-Mass Binary Systems

Sp. Type	Binary Separation Distance ($2a$)								
...	0.2	0.5	1.0	2.0	3.0	4.0	5.0	7.5	10.0
...	(AU)	(AU)	(AU)	(AU)	(AU)	(AU)	(AU)	(AU)	(AU)
F0	P	P	P	...	...	...	...	...	...
F1	P	P	P	...	...	...	...	...	...
F2	P	P	P	...	...	...	...	...	...
F3	P	P	P	...	...	...	...	...	...
F4	P	P	P	...	...	...	...	...	...
F5	P	P	P	...	...	...	...	...	...
F6	P	P	PT	...	...	...	...	...	...
F7	P	P	PT	...	...	...	...	...	...
F8	P	P	PT	...	...	...	...	...	...
F9	P	P	...	...	...	...	...	...	ST
G0	P	P	...	...	...	...	...	...	ST
G1	P	P	...	...	...	...	...	...	ST
G2	P	P	...	...	...	...	...	...	ST
G3	P	P	...	...	...	...	...	...	ST
G4	P	P	...	...	...	...	...	ST	ST
G5	P	P	...	...	...	...	...	ST	ST
G6	P	P	...	...	...	...	...	ST	S
G7	P	P	...	...	...	...	...	ST	S
G8	P	PT	...	...	...	...	...	ST	S
G9	P	PT	...	...	...	...	...	ST	S
K0	P	PT	...	...	...	...	...	ST	S
K1	P	PT	...	...	...	...	ST	S	S
K2	P	...	...	...	...	...	ST	S	S
K3	P	...	...	...	...	...	ST	S	S
K4	P	...	...	...	...	ST	ST	S	S
K5	P	...	...	...	...	ST	S	S	S
K6	P	...	...	...	ST	ST	S	S	S
K7	PT	...	...	...	ST	S	S	S	S
K8	PT	...	...	...	ST	S	S	S	S

Table 8—Continued

Sp. Type	Binary Separation Distance (2a)								
...	0.2 (AU)	0.5 (AU)	1.0 (AU)	2.0 (AU)	3.0 (AU)	4.0 (AU)	5.0 (AU)	7.5 (AU)	10.0 (AU)
K9	PT	...	...	ST	ST	S	S	S	S
M0	PT	...	...	ST	S	S	S	S	S

Note. — S and P indicate S-type and P-type habitability of the binary system, respectively, given by the full spatial extent of the radiative habitable zones. ST and PT indicate S-type and P-type habitability of the binary system, respectively, that is truncated by the additional constraint of orbital stability for a putative system planet.

Table 9. Assessment of the General Habitable Zone of Equal-Mass Binary Systems

Sp. Type	Binary Separation Distance ($2a$)								
... ...	0.2 (AU)	0.5 (AU)	1.0 (AU)	2.0 (AU)	3.0 (AU)	4.0 (AU)	5.0 (AU)	7.5 (AU)	10.0 (AU)
F0	P	P	P	PT	...	...	...	...	...
F1	P	P	P	...	...	...	...	...	...
F2	P	P	P	...	...	...	...	...	...
F3	P	P	P	...	...	...	...	...	...
F4	P	P	PT	...	...	...	...	...	...
F5	P	P	PT	...	...	...	...	...	...
F6	P	P	PT	...	...	...	...	...	...
F7	P	P	PT	...	...	...	...	...	ST
F8	P	P	PT	...	...	...	...	...	ST
F9	P	P	PT	...	...	...	...	...	ST
G0	P	P	PT	...	...	...	...	...	ST
G1	P	P	PT	...	...	...	...	...	ST
G2	P	P	PT	...	...	...	...	ST	ST
G3	P	P	...	...	...	...	...	ST	ST
G4	P	P	...	...	...	...	...	ST	ST
G5	P	PT	...	...	...	...	...	ST	ST
G6	P	PT	...	...	...	...	...	ST	ST
G7	P	PT	...	...	...	...	...	ST	ST
G8	P	PT	...	...	...	...	...	ST	ST
G9	P	PT	...	...	...	...	...	ST	ST
K0	P	PT	...	...	...	...	ST	ST	S
K1	P	PT	...	...	...	...	ST	ST	S
K2	P	PT	...	...	...	ST	ST	ST	S
K3	P	PT	...	...	...	ST	ST	ST	S
K4	P	...	...	...	...	ST	ST	S	S
K5	P	...	...	...	ST	ST	ST	S	S
K6	PT	...	...	...	ST	ST	S	S	S
K7	PT	...	...	...	ST	ST	S	S	S
K8	PT	...	...	ST	ST	S	S	S	S

Table 9—Continued

Sp. Type	Binary Separation Distance ($2a$)								
... ...	0.2 (AU)	0.5 (AU)	1.0 (AU)	2.0 (AU)	3.0 (AU)	4.0 (AU)	5.0 (AU)	7.5 (AU)	10.0 (AU)
K9	PT	...	...	ST	ST	S	S	S	S
M0	PT	...	...	ST	ST	S	S	S	S

Note. — See Table 8 for comments.

Table 10. Assessment of the Extended Habitable Zone of Equal-Mass Binary Systems

Sp. Type	Binary Separation Distance ($2a$)								
... ...	0.2 (AU)	0.5 (AU)	1.0 (AU)	2.0 (AU)	3.0 (AU)	4.0 (AU)	5.0 (AU)	7.5 (AU)	10.0 (AU)
F0	P	P	P	PT	PT	...	...	...	...
F1	P	P	P	PT	...	...	...	...	...
F2	P	P	P	PT	...	...	...	...	...
F3	P	P	P	PT	...	...	...	...	...
F4	P	P	PT	PT	...	...	...	...	...
F5	P	P	PT	PT	...	...	...	...	...
F6	P	P	PT	PT	...	...	...	...	...
F7	P	P	PT	PT	...	...	...	...	ST
F8	P	P	PT	...	...	...	...	...	ST
F9	P	P	PT	...	...	...	...	...	ST
G0	P	P	PT	...	...	...	...	...	ST
G1	P	P	PT	...	...	...	...	...	ST
G2	P	P	PT	...	...	...	...	ST	ST
G3	P	P	PT	...	...	...	...	ST	ST
G4	P	P	PT	...	...	...	...	ST	ST
G5	P	PT	PT	...	...	...	...	ST	ST
G6	P	PT	PT	...	...	...	...	ST	ST
G7	P	PT	PT	...	...	...	...	ST	ST
G8	P	PT	PT	...	...	...	...	ST	ST
G9	P	PT	PT	...	...	...	...	ST	ST
K0	P	PT	...	...	...	...	ST	ST	ST
K1	P	PT	...	...	...	...	ST	ST	ST
K2	P	PT	...	...	...	ST	ST	ST	ST
K3	P	PT	...	...	...	ST	ST	ST	ST
K4	P	PT	...	...	...	ST	ST	ST	S
K5	P	PT	...	...	ST	ST	ST	ST	S
K6	PT	...	...	...	ST	ST	ST	S	S
K7	PT	...	...	...	ST	ST	ST	S	S
K8	PT	...	...	ST	ST	ST	ST	S	S

Table 10—Continued

Sp. Type	Binary Separation Distance ($2a$)								
...	0.2 (AU)	0.5 (AU)	1.0 (AU)	2.0 (AU)	3.0 (AU)	4.0 (AU)	5.0 (AU)	7.5 (AU)	10.0 (AU)
K9	PT	...	...	ST	ST	ST	S	S	S
M0	PT	...	...	ST	ST	ST	S	S	S

Note. — See Table 8 for comments.

Table 11. Limits of habitability in a G2 V equal-star binary system

HZ Type	P	PT	ST	S
...	(AU)	(AU)	(AU)	(AU)
CHZ	0.6	0.8	8.3	12.0
GHZ	0.5	1.0	7.4	14.6
EHZ	0.5	1.5	7.4	21.0

Note. — The limits of P and PT constitute upper limits, whereas limits of S and ST constitute lower limits. No habitability is found between the limits PT and ST.